This Page Is Inserted by IFW Operations and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

As rescanning documents will not correct images, please do not report the images to the Image Problem Mailbox.

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 7:

C12N 15/12, 15/57, 15/67, 15/85, 9/64, C07K 14/72, C12Q 1/68, A61K 48/00

A1

(11) International Publication Number:

WO 00/49147

(43) International Publication Date:

24 August 2000 (24.08.00)

(21) International Application Number:

PCT/EP00/01368

(22) International Filing Date:

18 February 2000 (18.02.00)

(30) Priority Data:

199 07 099.7 60/120,848

19 February 1999 (19.02.99) DE

19 February 1999 (19.02.99) US

(71) Applicant (for all designated States except US): THER AGENE BIOMEDICAL LABORATORIES GMBH [DE/DE]; Am Klopferspitz 19, D-82152 Martinsried (DE).

(72) Inventor; and

(75) Inventor/Applicant (for US only): HAUSER-FUNKE, Charlotte [DE/DE]; Romanstr. 95, D-80369 München (DE).

(74) Agents: HELBING, Jörg et al.; von Kreisler Selting Werner, Deichmannhaus am Dom, D-50667 Köln (DE).


(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published

With international search report.

Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.

(54) Title: HORMONE-HORMONE RECEPTOR COMPLEXES AND NUCLEIC ACID CONSTRUCTS AND THEIR USE IN GENE THERAPY



(57) Abstract

The invention relates to the use of a nucleic acid contruct comprising at least one hormone responsive element and a transgene for preparing an agent for gene transfer. It further relates to particular nucleic acid contructs comprising at least one hormone responsive element and a transgene, wherein one of said at least one hormone responsive elements is not functionally linked to the transgene, vectors comprising such nucleic acid contructs and compositions of matter comprising such nucleic acid constructs wherein the hormone responsive elements of the constructs are coupled to a hormone-hormone receptor complex. The nucleic acid constructs, plasmids, and compositions of matter of the invention have applications in gene therapy, particularly in the treatment of human blood clotting disorders, such as hemophilia. They may also be used to up- or down-regulate target genes and for the delivery of vaccines.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Annenia	FI	Finland	LT	Lithuania	- SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	· SN	Senegal
ΑU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
ΑZ	Azerbaijan	GB	United Kingdom	MC	Мопасо	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB ·	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	15	Iceland	MW	Malawi	US	United States of America
CA	Canada	IТ	Italy	MX	Mexico	UZ.	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	zw	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand	2.,	Zimoaowe
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

WO 00/49147 PCT/EP00/01368

Hormone –Hormone Receptor Complexes and Nucleic Acid Constructs and Their Use in Gene Therapy

Background of the Invention

1. Object of the Invention

The invention relates to the use of a nucleic acid construct comprising at least one hormone responsive element and a transgene for preparing an agent for gene transfer. It further relates to particular nucleic acid constructs comprising at least one hormone responsive element and a transgene, wherein one of said at least one hormone responsive elements is not functionally linked to the transgene, vectors comprising such nucleic acid constructs and compositions of matter comprising such nucleic acid constructs wherein the hormone responsive elements of the constructs are coupled to a hormone-hormone receptor complex. The nucleic acid constructs, plasmids, and compositions of matter of the invention have applications in gene therapy, particularly in the treatment of human blood clotting disorders, such as hemophilia. They may also be used to up- or down-regulate target genes and for the delivery of vaccines.

2. Summary of the Related Art

Gene therapy is a method that holds great promise for many diseases and disorders. In general, it involves the transfer of recombinant genes or transgenes into somatic cells to replace proteins with a genetic defect or to interfere with the pathological process of

5

10

15

20

10

15

20

25

30

an illness. In principle, gene therapy is a simple method. In practice, many disadvantages must still be overcome.

Research in gene therapy has concentrated on ways to most effectively incorporate DNA into cells of a patient. Viral vectors are currently the widely used vehicles in clinical gene therapy approaches. In terms of efficacy in gene expression, the viral delivery systems have major advantages over techniques using DNA-lipid formulations as delivery vehicles or over mechanical methods, such as the gene gun. Although there are a variety of viral systems tested for gene therapeutical strategies, retroviral vectors and adenoviral vectors are presently the most widely used vehicles (Salmons, B. and Gunzburg, W. H., Hum. Gene Ther., Vol. 4, 129, 1993; Kasahara, N. A., et al., Science, Vol. 266, 1373, 1994; Ali, M., et al., Gene Ther., Vol. 1, 367, 1994.). Still, these systems have major disadvantages, such as potential viral contamination. Other safety concerns continue to hamper the development of clinical application of gene therapy using these viral systems. For example, recombinant retroviruses have the disadvantage of random chromosomal integration, which may lead to activation of oncogenes or inactivation of tumor-suppressor genes. Also, repetitive use of recombinant adenoviruses has caused severe immunological problems (Elkon, K. B. et al., Proc. Natl. Acad. Sci. USA, Vol. 94, 9814, 1997). The humoral response resulted in the production of antibodies to adenovirus proteins preventing subsequent infection. Immunosuppressive drugs may ameliorate these effects, but they place an additional burden on the patient (Dai, Y., et al., Proc. Natl. Acad. Sci. USA, Vol. 92, 1401, 1995).

Yet another viral delivery system involves adenoassociated virus (AAV). The AAV requires coinfection with an unrelated helper virus. Although such recombinant AAV virions have proven useful for introducing several small gene sequences into host cells, gene

10

15

20

25

30

delivery systems based on those particles are limited by the relative small size of AAV particles. This feature greatly reduces the range of appropriate gene protocols. Moreover, the need to also use a helper virus adds a complicating factor to this delivery system (Muzyczka, N., Curr. Top. Microbiol. Immunol., Vol. 158, 97, 1992).

Though safer, non-viral gene therapy approaches are also unsatisfactory. Problems with inefficient gene delivery or poor sustained expression are major drawbacks. Yet the methods available such as the direct injection of DNA into cellular compartments, or the application of mixtures of DNA with cationic lipids or polylysine allowing the transgene to cross the cell membrane more easily, have not overcome these hurdles (Felgner, P., et al., *Proc. Natl. Acad. Sci. USA*, Vol. 84, 7413, 1987; Behr, J.-P., *Bioconjugate Chemistry*, Vol. 5, 382, 1994).

Introduction of naked DNA (polynucleotide) sequences (including antisense DNA) into vertebrates, is reported to be achieved by injection into tissues such as muscle, brain or skin or by introduction into the blood circulation (Wolff, J. A., et al., Science, Vol. 247, 1990; Lin, H., et al., Circulation, Vol. 82, 2217, 1990; Schwartz, B., et al., Gene Ther., Vol. 3, 405, 1996). Also, a direct gene transfer into mammals has been reported for formulations of DNA encapsulated in liposomes and DNA entrapped in proteoliposomes containing receptor proteins. Although injected naked DNA leads to transgene expression. the efficiency is by far not comparable to viral-based DNA delivery systems. A limitation of the method of naked DNA injection is the fact that transgene expression is dose-dependent. The gene expression is saturable, and an increase in the amount of DNA injected leads to decreased protein production per plasmid. Thus, protein expression can dramatically decrease, if the amount of DNA injected is above a certain threshold.

10

15

20

25

30

Among the genetic disorders that the skilled artisan has sought to overcome using these prior art methods are those relating to blood clotting disorders, and in particular, hemophilia (Lozier, J. N. and Brinkhous, K. M., *JAMA*, Vol.271, 1994; Hoeben, R. C., *Biologicals*, Vol. 23, 27, 1995). For example, hemophilia A and B are X-linked, recessive bleeding disorders caused by deficiencies of clotting factors VIII and IX, respectively (Sadler, J. E. et al., in: *The Molecular Basis of Blood Diseases*, 575, 1987). The incidence of hemophilia is about 1 in 5,000 male births. Hemophiliacs suffer from excessive bleeding due to the lack of clotting at the site of wounds. The inability to clot properly causes damage to joints and internal tissues as well as posing risks to the proper treatment of cuts.

Treatment of hemophilia A is possible by the administration of the blood clotting factor VIII. Until recently, factor VIII preparations had to be prepared by concentrating blood from donors, posing the risk of contamination by infectious agents, such as HIV and hepatitis. The gene for factor VIII has been cloned (e.g., Vehar et al., *Nature* Vol. 312, 337 1984) allowing for the production of a recombinant product. Although recombinant methods provide factor VIII of higher purity than blood concentrates, the exogenous supply of factor VIII to a patient still requires repeated doses throughout the lifetime of the patient, an inconvenient and expensive solution.

Other forms of hemophilia include hemophilia B, caused by a defect in the gene coding for Factor IX. The gene therapy systems described above have been attempted for the treatment of hemophilia A and B with factors VIII and IX, respectively. (See e.g., WO 94/29471). However, these systems have the disadvantages already discussed above.

On the other hand, the classical model of the action of hormones is based on the concept of binding interaction of the hormone to an

15

20

25

intracellular receptor, located in the cytoplasm or the nucleus (Evans, R., *Science*, Vol. 240, 889, 1988). These intracellular receptors remain latent until exposed to their target hormone. When so exposed, the hormone receptor changes its conformation after the hormone is bound and translocates in the activated form into the cell nucleus where it binds as a dimer to hormone responsive elements in the promoter region of hormone-regulated genes (Beato, M., *Cell*, Vol. 56, 335, 1989; O'Malley, B., et al., *Biol. Reprod.*, Vol. 46, 163, 1992). The hormone responsive elements are enhancer elements usually located in the 5' flanking region of the specific hormone-induced gene, i.e., are functionally linked to the specific hormone induced gene. DNA constructs comprising a hormone responsive element and a nucleic acid sequence encoding a protein of interest are disclosed in U.S. Pat. Nos. 5,688,677 and 5,580,722 and are taught to be suitable for expression of the protein of interest.

An example of such intracellular receptors is the steroid receptor. Steroid receptors belong to a superfamily of ligand-dependent transcription factors characterized by a unique molecular structure. The centrally located highly conserved DNA-binding domain defines this superfamily. The second important and relatively invariant region is the COOH-terminal ligand-binding domain. An example of such a receptor is the progesterone receptor mediated by the steroid progesterone. At the progesterone receptor, progesterone acts as a natural agonist whereas it displays potent antimineralocorticoid properties both at the molecular and the systemic level. Besides classical effects on the uterus, antiepileptic, anxiolytic, hypnotic and anesthetic properties have been attributed to progesterone according to numerous studies.

Methods have been proposed for the use of mutant hormone receptors, including mutant steroid receptors for gene therapy. For

example, such methods are disclosed in WO 93/23431, WO 98/18925, WO 96/40911. Moreover, WO 98/33903 discloses a genetic construct comprising a steroid responsive element from a tissue specific gene, a coding sequence, and an SV40 enhancer.

5

10

15

20

25

Brief Description of the Invention

The object of the present invention is to overcome the disadvantages of the previous gene therapy delivery systems. It was found that a hormone-hormone receptor complex possesses the ability to drag a nucleic acid construct having one or more hormone responsive element(s) through the cell membrane into a cell. It was also found that if the construct comprises further functional sequences besides the hormone responsive elements (hereinafter "transgenes"), the functional sequences exert their function. The hormone responsive element may also enhance the expression of the transgene. Moreover, it was found that steroid hormones are very effective mediators for the transfer of nucleic acid constructs through the cell membranes into a cell. The present invention thus provides

- (1) the use of a nucleic acid construct comprising at least one hormone responsive element (hereinafter referred to as "HRE") and a transgene for preparing an agent for gene transfer (said at least one HRE being functionally linked to the transgene or not);
- (2) a preferred embodiment of (1) above, wherein the agent further comprises a hormone-hormone receptor complex;
- (3) a nucleic acid construct comprising at least one HRE and a transgene, wherein one of said at least one HREs is not functionally linked to the transgene;
 - (4) a vector comprising the nucleic acid construct of (3) above;

10

15

20

- (5) a transformed cell or transgenic organism comprising the nucleic acid construct as defined in (3) above or the vector as defined in (4) above;
- (6) a composition of matter comprising a nucleic acid construct comprising at least one HRE and a transgene as defined in (3) above and/or a vector as defined in (4) above, said at least one HRE being coupled to a hormone-hormone receptor complex;
- (7) a preferred embodiment of (6) above, wherein the transgene is a gene encoding a blood clotting factor;
- (8) a preferred embodiment of (7) above, wherein the blood clotting factor is factor IX;
- (9) a preferred embodiment of (7) above, wherein the blood clotting factor is factor VIII;
- (10) a pharmaceutical composition comprising the nucleic acid construct as defined in (3) above and/or the composition of matter as defined in (6) to (9) above;
- (11) a method for preparing the composition of matter as defined in (6) above, which method comprises admixing the nucleic acid construct with the hormone receptor and the hormone;
- (12) a method for gene transfer which comprises administering the agent as defined in (1) and (2) or the composition of matter as defined in (6) to (9) above to an organism or to a cellular system;
- (13) a method for delivering into an organism or into a cellular system a nucleic acid encoding a transgene to be expressed in the cells of the organism or the cells of the cellular system, which method comprises administering an agent as defined in (1) above or composition of matter as defined in (6) to (9) above to the organism or to the cellular system so that the hormone in the composition interacts with the cell membrane and therewith enhances diffusion

10

15

20

25

and transport of the nucleic acid that is coupled to the hormonehormone receptor complex across the membrane and into the cell;

- (14) a method of treating blood clotting disorders comprising administering a therapeutically effective amount of the composition of matter as defined in (7) above to an organism or to a cellular system;
- (15) a method of treating hemophilia B, comprising administering a therapeutically effective amount of the composition of matter as defined in (8) above to an organism or to a cellular system;
- (16) method of treating hemophilia A, comprising administering a therapeutically effective amount of the composition of matter as defined in (9) above to an organism or to a cellular system;
- (17) use of a steroid hormone for preparing an agent for gene transfer; and
- (18) a method for gene transfer which comprises administering a nucleic acid construct to an organism or to a cellular system, wherein the nucleic acid construct contains a transgene and is encapsulated in a steroid hormone.

In a preferred embodiment of (1) to (16) above the hormone responsive element is a steroid responsive element (SRE), most preferably a progesterone responsive element (PRE). In embodiments (2) and (6) to (16) the receptor preferably is a steroid receptor, most preferably, a progesterone receptor. Similarly, the hormone is preferably a steroid, most preferably, progesterone.

The present invention thus provides a delivery system for gene therapy that should overcome the prior art disadvantages. The presence of the hormone responsive element on the nucleic acid carrying a transgene encourages the binding of a hormone-hormone receptor complex. Thus, the present invention uses the activated hormone receptor as a link (or binding compound) between the

10

15

20

25

nucleic acid carrying the transgene and the hormone known to interact with the cell membrane. The general known biological activity mediated by the HREs is not the primary effect utilized in the present invention, but might the an additional effect when regulation of the transgene is desired. The general principle is depicted in Figure 1. The hormone responsive element is preferably present as a nucleic acid dimer sequence or nucleic acid multimer sequence. Even in an inverse orientation, the hormone responsive element will exert its proper function. The hormone-hormone receptor complex contains a hormone receptor that becomes activated after binding of its specific hormone. The hormone receptor in the activated state is able to recognize and bind to its specific hormone responsive element, which in the present invention is present within the nucleic acid comprising the desired transgene, e.g., a human blood-clotting factor.

Vaccination is another aspect of the embodiment (12) defined above. Introducing a nucleic acid construct or composition of matter of the invention comprising a gene for an antigen or containing a viral sequence into a cell (DNA vaccines) using the method mentioned above may also provide a way to stimulate the cellular immune response.

Brief Description of the Drawings

Figure 1 shows the concept of gene transfer of the present invention (with HRE = hormone responsive element, HR = hormone receptor, H = hormone, blank circles = lipophilic matrix).

Figure 2 is a diagram of the vector pTGFG1.

<u>Figure 3</u> is a diagram of the vector pTGFG5.

Figure 4 is a diagram of the vector pTGFG20.

Figure 5 is a diagram of the vector pTGFG33.

Figure 6 is a diagram of the vector pTGFG36.

Figure 7 is a diagram of the vector pTGFG53.

Figure 8 is a diagram of the vector pTGFG64.

Figure 9 is the DNA sequence of vector pTGFG36 (SEQ ID NO: 1).

Figure 10 shows the protein sequence of factor IX encoded by vector pTGFG36 (SEQ ID NO: 2).

<u>Figure 11</u> shows a GFP concentration curve for cell homogenates after transfection with pTGFG5 and pTGFG20, respectively.

Figure 12 shows corresponding light (a and c) and fluorescent (b and d) micrographs of HeLa cells transfected with pTGFG5 (a and b) and pTGFG20 (c and d), respectively.

Figure 13 shows the amount of GFP expressed by utilizing the favoured vectors of the invention in a transfection experiment.

Relative fluorescence units from mock and background can be clearly separated.

<u>Figure 14</u> shows the additive effect of human clotting factor IX on clotting activity of mouse blood.

Figure 15: hPR (A-form) was expressed in insect cells and purified by cobalt²⁺ affinity chromatography as described in Example 5. The final preparation (85µg protein) was separated on a denaturing 7,5% SDS-polyacrylamid gel, followed by staining with coomassie® R250 (lane A) or western blotting with hPR-specific staining (lane C).

Lane B: Molecular mass standard. Arrows indicate the two highly enriched protein species (94 and 74 kDa) accessible to

25 immunodetection.

Figure 16: Domain structure of hPR-B (numbers on the top of the bar represent amino acid positions within the polypeptide sequence).

Figure 17 shows the mean values of the difference in the clotting time of Example 9.

Figure 18 shows the clotting time detected in Example 9.

15

10

20

<u>Figure 19</u> shows the activity of human progesterone receptor as determined in Example 8.

<u>Figure 20:</u> shows the amino acid sequence of the hPR B-Form. The start methionine 165 of the hPR A-Form is underlined (SEQ ID NO: 18).

Figure 21 shows the nucleic acid sequence of the mRNA coding for hPR. The reading frame for the hPR B-form starts at position 176, the reading frame for the hPR A-Form at position 668. The respective start codons ATG are underlined (SEQ ID NO: 19). The sequences of Figures 20 and 21 are taken from Genbank, accession number AF016381.

Detailed Description of the Invention

1. Definitions

"Nucleic acid" means DNA, cDNA, mRNA, tRNA, rRNA. The nucleic acid may be linear or circular, double-stranded or single-stranded.

"Nucleic acid construct" refers to a composite of nucleic acid elements in relation to one another. The nucleic acid elements of the construct may be incorporated into a vector in such an orientation that a desired gene may be transcribed, and if desired, a desired protein may be expressed.

"Transgene" refers to a functional nucleic acid sequence which is transcriptionally active (with or without regulatory sequences).

"Gene transfer" includes "gene therapy".

"Hormone responsive element" (HRE) refers to regions of nucleic acids, and in particular, DNA, which regulate transcription of genes in response to hormone activation. HREs are typically about 10-40 nucleotides in length, and more usually, about 13-20 nucleotides in

5

10

15

20

25

length. As explained above, HREs become activated when a hormone binds to its corresponding intracellular receptor causing conformational change, so that the receptor has increased affinity for the HRE and binds to it. The HRE, in turn, stimulates transcription. A "steroid responsive element" (SRE) is an HRE that regulates transcription of genes in response to steroid activation. "progesterone responsive element" (PRE) is an HRE/SRE that regulates transcription of genes in response to progesterone activation.

A "hormone receptor" refers to a receptor which binds to and is activated by a hormone. A "steroid receptor" refers to a receptor which binds to and is activated by a steroid hormone. A "progesterone receptor" is a receptor which binds to or is activated by the steroid hormone progesterone.

"Functionally linked" refers to configurations of the nucleic acid construct, where the HRE (or SRE/or PRE) is located within the construct so that it can stimulate transcription of the transgene. "Not functionally linked" refers to configurations where the HRE is so remotely located from the transgene that it cannot stimulate its transcription.

"Gene" refers to DNA sequence encoding a polypeptide, optionally including leader and trailer sequences and introns and exons.

"Vector" refers to any genetic construct, such as a plasmid, phage, cosmid, etc., which is capable of replication when associated with the proper control elements and which can transfer gene sequences between cells. The term includes cloning and expression vehicles.

"Promoter" refers to a region of regulatory DNA sequences for the control of transcription of a gene to which RNA polymerase binds.

10

15

20

25

The promoter forms an initiation complex with RNA polymerase to initiate and drive transcription activity. "Enhancers" may activate the complex or "silencers" may inhibit the complex. A "tissue-specific promoter" is a promoter found in the DNA of tissue for transcription of genes expressed in this specific tissue.

"Organism" refers to a multicellular living entity including vertebrates such as mammals (especially humans, cattle, rodents, dogs) and invertebrates.

"Cellular system" includes cell cultures, e.g., primary cell cultures (especially those suitable for reimplantation), stem cells, blood cells, tissue samples and whole organs and immortalized cell cultures.

"Therapeutically effective dose" of the products of the invention refers to a dose effective for treatment or prophylaxis, for example, a dose that yields effective treatment or reduction of the symptoms of hemophilia. It is also a dose that measurably activates expression of a target gene as determined by measurements of target protein levels, or a dose that is predictable to be effective for treatment or prophylaxis by extrapolating from *in vitro* or *in vivo* data. The determination of a therapeutically effective dose is within the purview of one skilled in the art.

"Encodes" or "encoding" refers to a property of the nucleic acid sequence of being transcribed (in case of DNA) or translated (in the case of mRNA) into a polypeptide *in vitro* or *in vivo* when placed under the control of appropriate regulatory sequences.

For the purposes of this application, "express", "expressing" or "expression" shall refer to transcription and translation of a gene encoding a protein.

5

10

15

20

25

2. Detailed Description and Examples

As stated above, an object of the present invention is to provide a new and improved delivery system for gene therapy. The invention thus provides nucleic acid constructs comprising at least one HRE and a transgene wherein one of said at least one HREs is not functionally linked to the transgene, and compositions of matter comprising such nucleic acid construct wherein said at least one HRE is coupled to a hormone-hormone receptor complex (embodiments (3) and (6) defined above). A preferred embodiment of the nucleic acid construct and of the composition of matter of the invention is one where the hormone responsive element is a steroid responsive element (SRE), and the receptor is a steroid receptor. Most preferably, the hormone responsive element is a progesterone responsive element (PRE), and the receptor is a progesterone receptor.

Potential HREs for use in the present invention have been previously described. For example, GREs (Scheidereit, C., et al., *Nature*, Vol. 304, 749, 1983; von der Ahe, D., et al., *Proc. Natl. Acad. Sci. USA*, Vol. 83, 2817, 1986), EREs or PREs (Chambon, P., et al., *Rec. Prog. Horm. Res.*, Vol., 40, 1, 1984; Klock, G., et al., *Nature*, Vol. 329, 734, 1987). As already stated above, the most preferred HRE for the invention is a PRE. Specifically, the preferred PRE is described in Example 1, i.e., is the double stranded DNA sequence comprised of SEQ ID NOs: 3 and 4. The nucleic acid for use in the invention comprises at least one hormone responsive element. Preferred is a nucleic acid comprising more than one HRE. For example, the nucleic acid may comprise three to ten, preferably three to five HREs. The most preferred embodiment is a nucleic acid comprising three to five PREs.

Potential hormone receptors for use in the present invention are, for example, estrogen receptors, mineralocorticoid receptors,

10

15

20

25

30

alucocorticoid receptors, retinoic acid receptors, androgen, calcitriol, thyroid hormone or progesterone receptors and orphan receptors. Such receptors have been previously described. (Green, S., et al., Nature, Vol. 320, 134, 1986; Green, G. L., et al., Science, Vol. 231, 1150, 1986; Arriza, J. L., et al., Science, Vol. 237, 268, 1987; Hollenberg, S. M., et al., Nature, Vol. 318, 635, 1985; Petkovitch, M., et al., Nature, Vol. 330, 444, 1987; Giguere, V., et al., Nature, Vol. 330, 624, 1987; Tilley, W., et al., Proc. Natl. Acad. Sci. USA, Vol. 86, 327, 1989; Baker, A. R., et al., Proc. Natl. Acad. Sci. USA, Vol. 85, 3294, 1988; Weinberger, C., et al., *Nature*, Vol. 324, 641, 1986; Sap. J., et al., *Nature*, Vol. 324, 635, 1086; Misrahi, M., et al., *Biochem*. Biophys. Res. Commun., Vol. 143, 740, 1987; Kastner, P., et al., Cell, Vol. 83, 859, 1995). These receptors may be from human or other mammalian sources, although human is preferred. Nucleotide and/or amino acid sequences of human steroid receptors are available in the GenBank: mineralocorticoid receptor: M16801; glucocorticoid receptor M10901; glucocorticoid receptor α_2 : U01351; glucocorticoid receptor β : M11050; retinoic acid receptor α : AF088888 (exon 1), AF088889 (exon 2), AF088890 (exon 3), AF088891 (exon 4). AF088892 (exon 5 and 6), AF088893 (exon 7), AF088894 (exon 8), AF088895 (exon 9 and complete cDNA); retinoic acid receptor v: M24857; androgen receptor: M27423 (exon 1), M27424 (exon 2), M27425 (exon 3), M27436 (exon 4), M27427 (exon 5), M27428 (exon 6), M27429 (exon 7), M27430 (exon 8); thyroid hormone receptor α_1 : M24748, thyroid hormone receptor α_2 : J03239; progesterone receptor: AF016381; somatotropin receptor: J00148; vitamin D receptor (calcitriol receptor): J03258.

The skilled person will understand that expression of the receptor proteins can be achieved by standard methods, e.g. via PCR-cloning of the known cDNAs from cDNA libraries and overexpression of

10

15

20

25

the corresponding proteins in suitable expression vectors, such as, for example, the vectors of the present invention, in suitable host cells, e.g., COS cells. Accordingly, subsequent purification of the cytosolic fraction can be achieved by routine methods such as affinity chromatography purification. For this purpose, various suitable antibodies against the desired receptor are commercially available. For example, polyclonal antibodies against the mouse progesterone receptor that have a sufficiently high cross-reactivity for the human protein are available from Dianova (Hamburg, Germany). Likewise, further purification can be achieved by standard methods, e.g., chromatographical methods such as ion-exchange chromatography and/or FPLC.

The most preferred receptor is the progesterone receptor. Preferably, the receptor is a human progesterone receptor. Such a human progesterone receptor (from T47D human breast cancer cells) is disclosed in US Patent No. 4,742,000, and cells expressing this receptor have been deposited (ATCC deposit number HTB, 133). As already described above, it would be routine to purify such a receptor from the cytosol using receptor specific antibodies. In addition, US Patent No. 4,742,000 discloses a method for purification of the human progesterone receptor using a specific steroid affinity resin (cf. Grandics et al., Endocrinology, Vol. 110, 1088, 1982).

Briefly, the cytosolic fraction of the T47D cells is passed over Sterogel, a commercial preparation of deoxycorticosterone coupled to Sepharose® 2B that selectively binds the progesterone receptor. After washing with loading buffer, the bound receptor is eluted with a buffer containing progesterone. The eluted steroid-receptor complex is then chromatographed on DEAE-Biogel and eluted stepwise with a buffer containing 0.2M NaCl. Subsequently, the bound progesterone can be

10

15

20

25

readily exchanged. As described above, further purification can be achieved by routine methods well-known to the skilled person.

An alternative method is disclosed in Example 5.

The structure of the hPR polypeptide is depicted in Fig. 16. The hPR polypeptide is composed of distinct structural domains. Naturally the human progesterone receptor (hPR) is expressed as two different sized proteins termed hPR-B (120 kDa) and hPR-A (94 kDa). HPR-A is a truncated but otherwise identical form of hPR-B, that is missing 165 the N-terminal amino acids (see Fig. 20, SEQ ID NO: 18). Both forms seems to be indistinguishable regarding their progesterone or DNA binding properties. In human cells the A and B forms of hPR are produced from the same gene by alternate initiation of translation at two different AUG start sites within the same RNA transcript. As it was reported earlier hPR-A and B can be expressed in Spodoptera (Sf9) cells as biological fully active polypeptides (Christensen et al., Mol. Endocrinol. 5, 1755ff (1991); Elliston et al., JBC 267, 5193-5198 (1992)).

The carboxyl terminus of the hPR polypeptide as shown in Fig. 16 comprises a progesterone binding domain (PBD) but also contains sequences responsible for the association with heat shock proteins and receptor dimerization. The hinge region provides a flexible link between the DNA-binding domain (DBD) and the PBD but is also thought to contain elements for receptor dimerization as well as nuclear localization. Binding of the hPR to its corresponding target sites at the chromosomal DNA (PREs, Progesterone Responsive Elements) is known to be mediated by the DBD. The remaining N-terminal trans-activation domain (TAD) consists of regions specific for the *in vivo* function of the hPR as a transcriptional gene activator.

Even though the N-terminus also seems to contribute directly to the homodimerization of hPR after progesterone binding, it has been

15

20

25

demonstrated that a fragment comprising only the hinge region and minimal C-terminal fragment to mediate the **PBD** was the progesterone dependent hPR-hPR-interaction (Tetel et al., Mol. Endocrinol. 11, 1114ff. (1997). It is believed that genetically engineered hPR polypeptides lacking either in part or completely the ·TAD (amino acids 1 to 556) might be expressed as structurally stable and fully soluble dimers in the presence of progesterone. Complexes with such a truncated hPR (provided that said truncated hPR exhibits DNA-binding activity as well as progesterone-binding activity) may functionally replace the complexes with the full length form of the described recombinant hPR-A or hPR-B proteins, since still mediating the contact between the plasmid DNA and the progesterone. Thus, the hPR in embodiments (2) and (6) to (16) of the invention preferably is a PR comprising nucleic acids 557 to 933 of natural hPR shown in SEQ ID NO: 18.

Effective expression of such a truncated version of hPR is possible in the baculovirus system but also in other eukaryotic expression systems, such as cultivated mammalian cells or yeast cells. Furthermore, also an *E. coli* overexpression strain is a possible system for the production of those polypeptides. In this case, the fusion of such a truncated hPR-version to a suitable polypeptide sequence, e.g. a histidine containing sequence or the GST (glutathion S- transferase) protein, might be helpful to overcome insolubility problems as well as to facilitate the isolation and purification of the expressed protein.

Mutated versions of these receptors and derivatives thereof, that still retain the function of the receptors to bind a ligand and thereby become activated and bind DNA and regulate transcription, may also be employed in the invention. Such derivative may be a chemical derivative, variant, chimera, hybrid, analog, or fusion.

20

25

The third component of the gene transfer system of the invention is the hormone. The hormone in the agent of embodiment 2 and in the composition of matter of embodiment (6) include synthetic and natural hormones, preferably steroid hormones, such as estrogen, testosterone, glucocorticoid, androgen, thyroid hormone, progesterone or derivatives thereof. These are widely available. Progesterone is most preferred. For example, natural micronized progesterone is the preferred progesterone from which has been marketed in France since 1980 under the trademark of UTROGESTAN® and is still available in Germany under the trademark UTROGEST®. Its properties are similar to the endogenous progesterone, in particular, it has antiestrogen, gestagen, slightly antiandrogen and antimineralocorticoid properties. The natural micronized progesterone in said marketed products is dispersed in a matrix as described hereinbelow.

The above micronized progesterone has advantages that make it a suitable carrier for genes or nucleic acid constructs to target cells. Specifically, the synergistic effect of the double process micronization and suspension in long-chain fatty acids residues of an oil results in increasing progesterone absorption. It has been demonstrated that after oral administration of 100 mg UTROGESTAN®, peak plasma progesterone levels were obtained after 1-4 hours in most cases (Padwick, M. L., et al., Fertil. Steril., Vol. 46, 402, 1986). Later on, the levels declined substantially, although they were still elevated at 12 hours. Even at 84 hours the levels were slightly higher than baseline. A U.S. kinetic study confirmed earlier demonstrating bioavailability the of oral micronized progesterone. They showed a peak effect at 2 hours followed by rapid decrease in plasma progesterone level (Simon, J. A., et al., Fertil., Steril., Vol., 60, 26, 1993).

15

20

25

30

A further advantage of using progesterone as a carrier is the low level of disadvantageous side effects. Orally administered progesterone adversely affects neither plasma lipids (Jensen, J. et al., Am. J. Obstet. Gynecol., Vol. 156, 66, 1987) nor carbohydrate metabolism (Mosnier-Pudar, H. et al., Arch. Mal. Coeur, Vol 84, 1111, 1991). Further, progesterone does not affect liver enzymes (ASAT, ALAT, AFOS), sex-hormone binding-globulin (SHBG) synthesis or HDLcholesterol levels at daily doses of 200 mg and 300 mg. Although the plasma levels of deoxycorticosterone may increase substantially during UTROGESTAN® treatment, there are strong indications that the mineralocorticoid effects of this progesterone metabolite completely counteracted by the anti-mineralocorticoid effects of progesterone itself. This is apparent from a comparative study (Corvol, P., et al., In: Progesterone and progestins. Raven Press, New York, 179, 1983) in which oral UTROGESTAN® was capable of antagonizing the mineralocorticoid effects of $9-\alpha$ -fluorohydrocortisone.

In the agent of embodiment (2) and in the composition of matter of embodiment (6) of the invention the molar ratio of HRE (or SRE/or PRE) within the nucleic acid construct to hormone receptor is preferably from 1:1 to 1:10, more preferably from 1:2 to 1:5. On the other hand, the molar ratio of hormone to hormone receptor is preferably at least 1000:1, more preferably at least 10000:1. Thus, the hormone is present in a large excess relative to the hormone receptor and the HRE, which is desirable in view of the ability of the hormones to transfer nucleic acid constructs through cell membranes.

The skilled artisan will appreciate that the agent of embodiments (1) and (2) and the pharmaceutical composition of embodiment (10) may contain other components capable of assisting in introducing the nucleic acid into a cell for the purpose of gene therapy (matrix compounds). Specifically, the agent and the composition, especially

10

15

20

25

the hormone component thereof, may contain the following matrix compounds: glucose and related compounds (such as D-sorbitol, Dmannitol); solubilizing adjuvants (such as alcohols, e.g., ethanol); polyhydric compounds such as glycerine, polyethylene glycol and polypropylene glycol; nonionic surface active compounds, ionic surface active compounds such as lecithin; oily compounds such as sesame oil, peanut oil soybean oil, corn oil, etc.; starches and their derivatives such as cyclodextrines and hydroxyalkylated starches; stabilizers such as human serum albumin, preservatives such as benzyl alcohol and phenol; and the like. The preferred matrix contains B-cyclodextrine, glycerine, lecithin and/or corn oil. For example, the pharmaceutical composition of hormone-hormone receptor nucleic acid complex of the invention may be provided orally to humans or animals as a gelatin capsule. Progesterone therein (preferably in micronized form) could be present in a concentration of 50 to 1000 mg, preferably 200 -300 mg dissolved in a 35 % or 40 % β-cyclodextrin solution or in cornoil or gycerol with peanut oil together with lecithin.

Alternatively, when - due to the selection of appropriate matrix components - the pharmaceutical composition is in a pasty, gel-like form, it may be provided topically.

The nucleic acid construct of embodiments (1) to (16) of the present invention may - aside from the transgene and the HREs, SREs, or PREs already disclosed above - further contain promoter, enhancer, and/or silencer sequences. The promoter may be ubiquitous or tissue-specific. Of the ubiquitous promoters, the CMV promoter is most preferred. However, a tissue-specific promoter is preferred over a ubiquitous promoter. For example, the tissue-specific promoters envisioned for the instant invention include α_1 -antitrypsin (further promoters).

15

20

25

30

The nucleic acid construct may further comprise additional sequences such as the ampicillin resistance gene. Other reporter sequences known to the skilled artisan may also be included, such as, for example, the green fluorescent protein (GFP), luciferase, ß-galactosidase or chloramphenicolacetyltransferase (CAT). As an enhancer sequence, the SV40 intron and SV40 Poly A are most preferred. The nucleic acid construct may further contain inducible promoters such as, for example, a MMTV (Mouse Mammary Tumor Virus) promoters inducible via glucocorticoides and Ecdyson-inducible insect promoters.

A preferred nucleic acid construct contains sequentially from the 5' to the 3' end: a PRE, a CMV promoter, a gene of interest, SV40 Intron and SV40 poly A enhancer sequence, and an ampicillin resistant gene. Further PREs are evenly distributed on the vector backbone.

The nucleic acid construct may further contain origin of replication sequences (especially eukariotic origin of replication sequences), elements for gene targeting, integrational sequences (e.g., AAV-ITR, transposon IS), 3'-UTR, "switch" systems (e.g., TET system, Cre/loxP or Flp/ftr system).

The transgene may be chosen from those encoding proteins lacking in a variety of genetic disorders or involved in conditions related to inappropriate responses to hormones, for example, hormone-dependent cancers such as breast, ovarian, and endometrial cancers and prostate cancer. The transgene may also be used to replace a defective gene resulting in such genetic disorders as hemophilia, von Willebrand disease, and cystic fibrosis. The transgene includes mutations of such gene or a gene encoding a fusion product. The nucleic acid construct of the present invention may comprise more than one transgene.

15

20

25

30

In particular, the transgene may replace genes for a blood clotting factor, and preferably a human blood-clotting factor. The genes encoding factor VIII and factor IX (sown in Fig. 2, SEQ ID NO: 2), involved in hemophilia A and B, respectively, are good candidates for the invention. Other candidates include the gene encoding von Willebrand factor, factor IV, factor X, or protein C.

Other useful transgenes include, but are not limited to, hormone genes such as the genes encoding for insulin, parathyroid hormone, luteinizing hormone releasing factor (LHRH), α and β seminal inhibins and human growth hormone; hormone receptor genes such as the glucocorticoid receptor, the estrogen receptor, the progesterone receptor, the retinoic acid receptor; growth factors such as vascular endothelial growth factor (VEGF), nerve growth factor, epidermal growth factor; enzyme genes; genes encoding cytokines or lymphokines such as interferons, granulocytic macrophage colony stimulating factor (GM-CSF), colony stimulating factor-1 (CSF-1), tumor necrosis factor (TNF), and erythropoietin (EPO); genes encoding inhibitor substances such as α_1 -antitrypsin, and genes encoding substances that function as drugs, e. g., genes encoding the diphteria and cholera toxins, ricin or cobra venom factor. Also, antisense sequences may be administered as genetic material.

Another aspect of the present invention is vectors comprising the nucleic acid constructs of embodiment (3) of the present invention. These vectors may be used in the composition matter of embodiment (6) of the present invention. Preferably, however, the nucleic acid sequence for use in the invention is circular rather than linear. The vectors may be capable of expressing the nucleic acid in the nucleic acid construct transiently or permanently (including episomally). As noted above, the nucleic acid construct therein may further contain additional elements.

15

20

25

The composition of matter of embodiment (6) of the invention can be prepared by admixing the nucleic acid construct with the hormone receptor and the hormone. Preferably, an aqueous solution of nucleic acid construct was added to the oily suspension containing the hormone at ambient temperature under stirring.

Embodiment(s) of the invention relates to transfected and transformed cells or transgenic organism comprising these vectors and/or nucleic acid constructs. Within the scope of this invention, a transfected cell is one in which foreign DNA has been incorporated. Methods of transfection may include microinjection, $CaPO_4$ precipitation, electroporation, liposome fusion, or gene gun.

Transformation refers to introducing genetic material into a cell, such as the vectors or nucleic acid constructs of the invention, rendering the cell transiently or permanently altered so that the cell expresses a specific gene product or is otherwise altered in its expression. Transformation may be achieved by *in vivo* or *in vitro* techniques, although *in vivo* transformation is preferred.

A further embodiment of the present invention is pharmaceutical compositions comprising a therapeutically effective dose of the nucleic acid constructs of the invention and a hormone. The hormone is preferably a steroid, and most preferably, progesterone, as described above. The dose is dependent on the condition to be treated, the characteristics of the patient, and the result sought to be achieved. Determining dosage is within the realm of the skilled artisan.

The pharmaceutical composition (or, alternatively, the composition of matter, the nucleic acid construct, or the vector) of the present invention may be administered orally, intravenously, intramuscularly, subcutaneously, topically, through mucosa (including buccal, nasal spray) or by gene gun. Oral administration (of a

10

20

25

micronized hormone dispersion) is preferred. Delivery may be systemic or directed at certain tissue.

The invention further includes a method of introducing into a cell a nucleic acid construct encoding a gene of interest, e.g., a human blood-clotting factor, to express the blood-clotting factor in the cell. In this method, the nucleic acid encoding a human blood-clotting factor is combined with a nucleic acid construct comprising at least one hormone responsive element (HRE), preferably a progesterone responsive element.

The mixture of nucleic acid bound to the hormone-hormone receptor complex together with an excess of hormone, preferably progesterone, will be used to introduce the nucleic acid into a cell by various methods known to the skilled artisan and outlined above. The cell-uptake will be stimulated by the interaction of the hormone with the cell membrane. The hormone or steroid interacts with the lipid bilayer of the cell membrane not only through membrane perturbation but also through activation of certain hormone- or steroid-sensitive membrane receptors. This has been demonstrated for progesterone and other steroids. Last but not least, it is known that hormones are able to cross the cell membrane by diffusion. In the present invention, the nucleic acid bound to the hormone-hormone receptor complex should be transported through the membrane during the process of diffusion or uptake.

Another aspect of the invention is a method of treating a blood clotting disorder by administering a therapeutically effective amount of the composition of matter of the invention to an organism. This method involves the administration and dosage considerations already discussed.

Embodiments (17) and (18) of the invention pertain to the use of a steroid hormone for preparing an agent for gene therapy and/or

gene transfer and to method for gene therapy and/or gene transfer which comprises administering a nucleic acid construct to an organism or to a cellular system, wherein the nucleic acid construct contains a transgene and is encapsulated in a steroid hormone. Suitable steroid hormones are enumerated hereinafter. The preferred steroid hormone in said embodiments of the invention is a natural micronized steroid hormone, in particular a natural micronized progesterone. In a preferred embodiment, the micronized hormone is solubilized/dispersed in a lipophilic matrix as described hereinafter.

Experiments have been performed to illustrate the technical aspects of the present invention. These experiments are described in examples 1 to 9 below. The skilled artisan will be readily recognize that the invention is not limited to these examples.

Examples

10

15

20

25

30

Example 1: Construction of Vectors

<u>Production of the vector pTGFG1:</u> The vector pUC19 (MBI Fermentas) was digested with XbaI, treated with Klenow enzyme and religated. This XbaI deleted vector was then digested with EcoRI, treated with Klenow enzyme and religated in order to delete the EcoRI site. For insertion of a XbaI site in the SacI site of this vector it was digested with SacI, treated with T4-polymerase, dephosphorylated with alkaline phosphatase and ligated with the XbaI-linker CTCTAGAG (Biolabs #1032). Another XbaI-site was inserted by digesting the newly produced vector with HindIII, treating it with Klenow, dephosphorylating it with alkaline phosphatase and ligating it with the XbaI-linker CTCTAGAG (Biolabs #1032). This vector was named pUC19/X.

15

20

In order to destroy the XbaI-site present in the vector phGFP-S65T (Clontech) this vector was digested with XbaI, treated with Klenow enzyme and religated resulting in the vector pGFP/0. A 2.3 kb fragment containing the GFP-Gene was isolated after digesting pGFP/0 with MluI, treating it with Klenow enzyme and digesting it with BamHI. This fragment was inserted into the multiple cloning site of the vector pUC19/X which was digested with SaII, treated with Klenow enzyme and digested with BamHI. The resulting vector was named pTGFG1 (Figure 2).

Starting with this vector all the vectors described in Table 1 were obtained. At the restriction sites for PstI, KpnI, Ehel, EcoO109 and/or SapI a PRE(ds) was inserted giving rise to plasmids carrying the GFP gene and up to five PREs. By exchanging the GFP gene with a FIX gene a set of FIX expression plasmids were obtained. By excising the GFP gene the cloning vectors without a transgene were obtained.

Production of the insert PRE(ds): The oligonucleotides (Metabion) PRE-S (5'-GGG GTA CCA GCT TCG TAG CTA GAA CAT CAT GTT CTG GGA TAT CAG CTT CGT AGC TAG AAC ATC ATG TTC TGG TAC CCC-3'; SEO ID NO: 3) and

PRE-AS (5'-GGG GTA CCA GAA CAT GAT GTT CTA GCT ACG AAG CTG ATA TCC CAG AAC ATG ATG TTC TAG CTA CGA AGC TGG TAC CCC-3'; SEQ ID NO: 4)

were hybridized and phosphorylated by kinase reaction, resulting in the insert PRE(ds).

<u>Production of the vector pTGFG5:</u> The vector pTGFG1 was digested with EcoO109I, treated with Klenow enzyme and dephosphorylated with alkaline phosphatase. It was then ligated with the PRE(ds) insert,

resulting in the vector pTGFG5 (Figure 3), i.e., a vector which carries a PRE at position C of Fig. 2.

Production of the vector pTGFG20: The vector pTGFG1 was digested with KpnI, treated with T4-polymerase and dephosphorylated with alkaline phosphatase. It was then ligated with the PRE(ds) insert, resulting in the vector pTGFG7. This vector pTGFG7 was digested with PstI, treated with T4-polymerase and dephosphorylated with alkaline phosphatase. It was then ligated with the PRE(ds) insert, resulting in the vector pTGFG11. Subsequently, pTGFG11 was digested with EcoO109I, treated with Klenow enzyme and dephosphorylated with alkaline phosphatase. It was then ligated with the PRE(ds) insert, resulting in the vector pTGFG20 (Figure 4). This vector carries a PRE at positions A, B and D of Fig. 2.

15

20

25

30

10

5

<u>Production of the vector pTGFG33:</u> In a similar manner PRE(ds) were inserted at the restriction sites for PstI, KpnI, EheI, EcoO109 and SapI in vector pTGFG1 giving rise to the plasmid pTGFG33 (Figure 5), which is a vector that carries the GFP gene and five PREs, one each in position A, B, C, D, E (Figure 2).

<u>Production of the vectors pTGFG36, pTGFG53 and pTGFG64:</u> The vector pUC19 (MBI Fermentas) was digested with SalI, treated with Klenow enzyme and dephosphorylated with alkaline phosphatase. It was ligated to the NotI-linker GCGGCCGC (Biolabs # 1045), resulting in the vector pUC19/N.

A 1.4 kb fragment containing the open reading frame of the human clotting factor IX, isolated from a human cDNA library (see example 2), was inserted into the PstI-site of the vector pUC19/N which was digested with PstI, treated with T4-polymerase and

15

20

dephosphorylated with alkaline phosphatase. From the resulting vector pUC19/N-FIX a 1.4 kb fragment containing the open reading frame of the human clotting factor IX was cut out by double-digestion with Hind III and NotI. This fragment was ligated to the 4.3 kb fragment of the HindIII and NotI double-digested vector pTGFG5 resulting in the vector pTGFG36 shown in Figure 6. This vector is a preferred one for delivery of Factor IX into the cell, and its DNA sequence is provided in Figure 9 (SEQ ID NO: 1).

In a similar manner plasmids pTGFG53 and pTGFG64 (shown in Figures 7 and 8) were obtained by exchanging the GFP gene in plasmids pTGFG20 and pTGFG33 by the FIX gene.

Production of the insert ALLG(ds): The oligonucleotides (Metabion) ALLG1/1 (5'-AGC TTG ACC TCG AGC AAG C-3') (SEQ. ID NO: 5) and ALLG2 (5'-GGC CGC TTG CTC GAG GTC A-3') (SEQ. ID NO: 6) were hybridized and phosphorylated by kinase reaction, resulting in the inserts ALLG(ds). The insert ALLG (ds) was constructed to introduce into the vector of choice a sequence with a multiple cloning site for the possible introduction of other transgenes.

Table 1 gives an overview of the available vectors with different transgenes and a different number of PREs in various positions. The positions of the PREs are given according to Figure 2. For the underlined vectors a map is provided (Figures 3 to 8).

Table 1: Vectors of the invention

Plasmid	Trans- gene	PRE	Plasmid	Trans- gene	PRE	Plasmid	Trans- gene	PRE
pTGFG0	-		pTGFG18	GFP	BDE	pTGFG34	FIX	E
pTGFG1	GFP		pTGFG19	GFP	BCD	pTGFG35	FIX	А
pTGFG2	FIX		pTGFG20	GFP	ABD	pTGFG36	FIX	D
pTGFG3	GFP	E	pTGFG21	GFP	CDE	pTGFG37	FIX	. C

		r						
pTGFG4	GFP	٨	pTGFG22	GFP	ACD	pTGFG38	FIX	В
pTGFG5	GEP	D	pTGFG23	GFP	ABC	pTGFG53	FIX	ABD
pTGFG6	GFP	С.	pTGFG24	GFP	ABE	pTGFG64	FIX	ABCDE
pTGFG7	GFP	В	pTGFG25	GFP	ACE	pTGFG66		Α
pTGFG8	GFP	ВС	pTGFG26	GFP	ADE	pTGFG67		D
pTGFG9	GFP	BE	pTGFG27	GFP	BCE	pTGFG68		C
pTGFG10	GFP	BD	pTGFG28	GFP	BCDE	pTGFG69		В
pTGFG11	GFP	AB	pTGFG29	GFP	ACDE	pTGFG82		ABD
pTGFG13	GFP	CD	pTGFG30	GFP	ABCE	pTGFG95		
pTGFG14	GFP	AC	pTGFG31	GFP	ABDE	p. G. G/3		ABCDE
pTGFG15	GFP	DE	pTGFG32	GFP	ABCD			
pTGFG16	GFP	AD	pTGFG33	GFP	ABCDE			
L								

For the DNA sequence of pTGFG 36, pTGFG 53, pTGFG 64, pTGFG 67, pTGFG 82 and pTGFG 95, see SEQ ID NOs: 1 and 13 to 17, respectively.

Example 2: Isolation of Human Factor IX cDNA

Factor IX cDNA was amplified from human liver cDNA (Clontech) using two primers overlapping the start and termination codon of the factor IX open reading frame resulting in a 1387 bp fragment containing the entire open reading frame. Restriction sites for EcoRI (upstream) and BamHI (downstream) were included at the end of each primer to facilitate cloning. Amplification was performed with Pwo polymerase (Boehringer Mannheim) in 50 ml reaction volume [10 mM Tris HCl pH 8.85, 25 mM KCl, 5 mM (NH₄)₂SO₄, 2 mM MgSO₄] with 30 incubation cycles at 96°C for 1 min, 60°C for 1 min, 72°C for 2 min, followed by a final extension step at 72°C for 10 min.

Reaction products were ligated into the EcoRI- and BamHI-sites of pUC19 and transformed into $\it E.~coli$ DH5-a. Positive clones were

selected. Sequences were confirmed by cycle sequencing (Amersham) from both ends with labeled primers (IR-700) and automated analysis on the LiCor sequencing system (MWG, Biotech).

The following primers were used:

GGAATTCCGCAAAGGTTATGCAGCGCGTGAACATGATCATGGC
(upstream; SEQ. ID NO: 7)
CGCGGATCCATTAAGTGAGCTTTGTTTTTCCTTAATCC (downstream;

SEQ. ID NO: 8)

10 Example 3: Expression and Quantification of the Marker Protein GFP

Method 1: HeLa cells were transfected by electroporation with plasmids pTGFG5 or pTGFG20. Transfected cells were harvested and the cell pellets were homogenized and lysed in a buffer containing phosphate buffered saline (pH 7.5) and 10 mM PMSF. The concentration of green fluorescent protein (GFP) in the cell homogenate was determined by competitive ELISA.

For this purpose, GFP was coated in a defined concentration on microtiter plates. Then, GFP samples were added in the presence of anti-GFP antibody. After several washing steps a labeled secondary antibody was added in order to trace the first antibody. The colorimetric reaction was measured photometrically (extinction). Generally, the more GFP was added the less antibody was left to bind the coated GFP. Thus, reduction of extinction corresponded to higher GFP concentration in the sample.

A concentration curve of GFP was determined by linear regression (Figure 11) using bovine serum albumin (BSA) as a reference. A mean value of 2.4 mg GFP/ml for pTGFG5 (1 PRE) and 5.2 mg GFP/ml for pTGFG20 (3PREs) was found.

15

20

25

Figures 12 a-d show micrographs of HeLa cell cultures transfected with pTGFG5 (Fig. 12 a and b) and pTGFG20 (Fig. 12 c and d), respectively. Figures 12 a and c represent light microscopic views as controls, and Fig. 12 b and d show the corresponding cell patches in the fluorescent mode. Routinely, more than 50% of the cells expressed GFP, indicating very efficient transfection, the presence of only one PRE showing more efficient expression.

Method 2: 293 T cells were transfected with pTGFG 5, 20 and 33 using calcium phosphate method and fluorescence was detected with a fluorimeter (Labsystems, Extinction: 485 nm Emission: 520 nm). In the case of the mock transfection, non GFP-expressing DNA was used. Background indicates the fluorescence of the empty plate (96-well plate, Dynex, Immulon-4). The results are summarized in Fig. 13.

15 Again the vector with just one PRE (pTGFG5) shows the highest expression.

Example 4: Human Factor IX Quantification by ELISA Assay

HeLa cells were transfected either by electroporation or using liposome reagent DOTAP (Boehringer Mannheim) with plasmids pTGFG36, pTGFG53 and pTGFG64. These plasmids contain the cDNA of human clotting factor IX. Recombinant human factor IX was secreted into the supernatant of the cell culture and quantified using a sandwich ELISA method.

0.11 M sodium citrate and 10 mM PMSF were added in order to prevent degradation of human factor IX. The enzyme-immunological in vitro assay "Asserachrom IX:AG" from Boehringer-Mannheim was used in order to determine the concentration of expressed human

clotting factor IX. The factor IX-standard from Octapharma AG was used as a standard in aqueous solutions of 28 IU/ml.

In six different transfection experiments, in which HeLa cells with plasmids containing human factor IX-cDNA (pTGFG36, 53 and 64) were transfected using either electroporation or lipid-transfection reagent (DOTAP, Boehringer Mannheim), a concentration range of 3-25 ng/ml human clotting factor IX was reached.

Example 5: Production and Purification of hPR (A Form)

10

- 1. Cloning of the human progesterone receptor: The cloning was performed as follows: Total human RNA was isolated from human white blood cells or liver cells using cell lysis in guanidinium hydrochloride buffer and CsCl-density centrifugation.
- For cloning of the hPR coding sequence, hPR specific cDNA was prepared and used for amplification of the hPR coding sequence in two fragments by PCR.
 - The following oligonucleotide primers were selected based on the published mRNA sequence (Genbank: NM_000926 and X51730).
 - Oligonucleotides used were obtained from MWG, Ebersberg or Metabion, München. All primers used are listed 5' to 3', bases added to introduce restriction sites are in capital letters and restriction sites used for cloning are underlined.
 - hPGR-5'-primer: CGA GGA tcc agt cgt cat gac tga gc (SEQ ID NO: 9);
- hPGR-3'-primer: GCA <u>GAA TT c</u>at tat aaa aac tca aga cct cat aat cct gac (SEQ ID NO: 10);
 - hPGR-internal primer (Sal I) 1: ctc ctc ggg gtc gac cct gg (SEQ ID NO: 11);
- hPGR-internal primer (Sal I) 2: cca ggg tcg acc ccg agg ag (SEQ ID NO: 12).

Synthesis of cDNA was perfomed using 3 µg of total RNA and 200 pmol of the 3'-primer with SuperScript II reverse transcriptase (Gibco BRL). Reaction volume was 50 µl and buffer was used as recommended, supplemented with RNase Inhibitor and 10 mM DTT and 1 mM dNTPs. Before adding the enzyme, samples were heated to 80°C for 10 min, followed by 10 min at 72°C and 10 min at 42°C. SuperScript II RT was added at 42°C and reaction was continued for 15 min at 42°C, 15 min at 50°C and 1 h at 58°C.

10

15

5

The cDNA obtained from this synthesis reaction was used to amplify the hPGR coding sequence in two fragments by PCR. One fragment (5') with 5'-primer and internal primer 2 and one fragment (3') with 3' primer and internal primer 1. Reaction setup in 50 µl was : Pwo polymerase (Roche Diagnostics), buffer as supplied by Roche Diagnostics, supplemented with DMSO, 50 pmol of each primer and 0.2 mM dNTPs. Reaction conditions were: 10 min 96°C followed by 35 cycles of 1 min 96°C, 2 min at 59°C, 2 min 72°C and a final extension step at 72°C for 10 min.

20

PCR-products were purified by gel electrophoresis and digested with Sal I. The BamHI and Hind III sites introduced in the primer were not used to avoid cutting at two internal restriction sites of the hPR coding sequence. Both fragments were ligated into pBluescript SK+ vector cut with EcoRV through blunt end ligation into the vector and sticky end ligation through the internal Sal I site. Vectors containing the appropriate insert were identified by mini-prep, restriction digest and sequencing. The obtained vector was designated pTGhPR1.

2. Production of hPR (A-form): Initially, the gene for hPR-B inclusive its 3'-UTR was cut out from pTGh PR1 and cloned in frame in the multiple cloning site of the expression plasmid pFASTBAC HTc (BAC-to-BAC Baculovirus Expression System, Life Technologies). This resulted in an expression casette of a N-terminally histidine-tagged version of hPR-B under expression control of the viral polyhedrin promotor as shown below. A rTEV protease cleavage site is located between the six histidine residues and the initial methionine of the hPR-B reading frame, which allows removal of the histidine residues from the expressed protein. The N-terminal region of the expression cassettes is shown below.

MSYY HHHHHH DYDIPTTE:	NLYF	Q**GP	MGIR	NST-h	PR-gen
6 x His					
spacer					*

rTEV cleavage site

Amino acids are presented in the single letter code. The cleavage site of the rTEV protease is represented by **

20

25

10

In order to generate the expression cassette for the truncated hPR-A form, the DNA sequence encoding for the amino acids between Met 1 and Met 165 of the hPR-B form was removed using a PCR-based strategy. Two primer pairs were designed which allowed amplification of either a DNA fragment just downstream of the start AUG of the hPR-B gene and a DNA-fragment just upstream of the AUG coding for Met 165, respectively. In a subsequent PCR reaction these two DNA fragments were annealed to each other at their homologous 3'-ends, and amplified using the outermost amplification primers. The resulting DNA-fragment was digested by EcoRI and Mlu I and the cleavage product was exchanged against the corresponding fragment of the

10

15

20

25

hPR-B expression cassette in the pFASTBAC HTc vector. Thereby the reading frame coding for an N-terminal histidine tagged version of the hPR-A polypeptide (94kDA) was restored.

This $6 \times \text{His-tag}$ was utilised for affinity purification of the protein by immobilized cobalt²⁺ affinity chromatography on a TALON® resin (Clontech). The procedure, following the method of Boonyaratanakornkit et al. Mol. Cell. Biol.18, 4471 (1998), was as follows (all steps were carried out at 0 to 8°C):

Sf9 cells were cultivated in monolayer culture in serum free SF900 medium. Viral infection of the cells was done at a multiplicity of infection (MOI) of 5-8.

The harvesting was done 48 hours after infection with baculovirus containing the hPR expression cassette and lysed mechanically by homogenising in buffer A containing 20 mM Tris-Cl pH 8.0, 350 mM NaCl, 10 mM imidazol, 5% glycerol and a cocktail of proteinase inhibitors (CompleteTM EDTA-free, Roche Diagnostics, Penzberg, Germany). After a 10 min centrifugation at $10000 \times g$, supernatant originating from 10^8 cells was incubated for 1 h with 0,5 ml settled TALON® resin equilibrated in buffer A. TALON® was washed with 20 volumes of buffer A. hPR-A was eluted with 10 Vol buffer B, containing all ingredients of buffer A, but 100 mM imidazol. The eluate was concentrated 50-fold and dialysed against 100 volumes buffer C (PBS + 100 nM progesteron) by centrifugal ultrafltration at a molecular exclusion size of 10 kDa (Centricon Plus-20 PL-10, Millipore, Eschborn, Germany).

3. Determination of identity, purity and yield of hPR-A: Purity and yield of the product were determined by application on denaturing reducing polyacrylamid- gelelectrophoresis according to Laemmli, U. et al., Nature 227, 680-685 (1970) and subsequent staining with

15

20

25

coomassie[©] blue R250. By this one-step procedure hPR-A was enriched to a final specific hPR content of 0.2 - 0.5 mg hPR/mg protein. As depicted in Figure 15, lane A, the final preparation consisted predominantly of two distinct protein species displaying apparent molecular masses of 94 and 74 kDa (Fig. 15, arrows).

Yield was estimated by parallel separation of standardised protein preparations. Data taken from a set of three separate experiments hint at a typical yield of 30 μ g enriched hPR A-receptor per 10^8 cells.

Identity of hPR was determined by immunodetection of the product transferred to nitrocellulose by western blotting with mouse monoclonal antibodies directed against recombinant hPR (PR Ab-1, Oncogene, Cambridge, MA, USA).

The final product was transferred to nitrocellulose BA-83 and immunostained as described above. As presented in Figure 15, lane C, three major protein bands were detected, including the two dominant protein species described above. The smaller sized bands may display copurified proteolytic fragments of hPR.

Intracellular GFP from adherent cells was detected by a fluorimeter after media was taken off and PBS (colourless) was added. The results are summarized in Fig. 13.

Example 6: Clotting Activity of Human Clotting Factor IX from Transfected 293 T Cells

A concentration range of 55 - 95 ng/ml human clotting factor IX has been reached by transfection of 293 T-cells with plasmids containing human factor IX-cDNA (pTGFG 36, 53, 64 and 2) in 11 different experiments using ELISA (Example 4).

Clotting activity was deterined with a partial thromboplastin time

assay using Cephalin (phosphatidyl ethanolamine) activation with a manual coagulation instrument (ML-2, Instrumentation Laboratories). For the study, 100 μ l undiluted supernatant from transfected 293 T-cells, 100 μ l deficiency plasma (Progen) and 100 μ l Cephalin (Instrumentation Laboratories) were incubated for 5 minutes at 37°C. Coagulation was started by adding 100 μ l CaCl₂. Sample coagulation time was compared to normal plasma.

Number of cells [/ml]	Factor IX-concentration [ng/ml]	Clotting time		
2.1×10^5	36	45		
$8,7 \times 10^5$	20	79		

10 Normal plasma:

37 - 39 s

Factor IX deficient plasma:

137 - 140 s

Example 7: Analysis of an Additive Effect of Human Clotting Factor IX on the Clotting Time of Mice Blood

1. Clotting time: Clotting activity was determined with a partial thromboplastin time assay using Cephalin (phosphatidyl ethanolamine) activation with a manual coagulation instrument (KC 4 A, Amelung).

For the study, 5 μ l mouse blood, 20 μ l deficiency plasma (Progen) ad 100 μ l physiological NaCl and 100 μ l DaPPTin (Progen) were incubated for 2 minutes at 37°C. Coagulation was started by adding 100 μ l CaCl₂.

To analyse the additive effect, human clotting factor IX (housestandard, Octapharma) was added to the mouse blood and diluted 1:10 within the system. As it is shown in Figure 15, the additive effect of human clotting factor IX on clotting activity can be

10

15

detected up to a limit concentration of 0.07 mIU hFIX/ml (= 31.5 ng/ml).

2. ELISA: The addition of human clotting factor IX to the mouse blood was monitored by ELISA as described in Example 4. Citrate plasma was made out of mouse blood and human clotting factor IX was added in different concentrations.

No.	Description	Concentration [mIU/ml] hFIX added	Extinction at 405 nm [-]			
1.	Mouse Citrate Plasma	7	0,204			
2.	Mouse Citrate Plasma	2	0,130			
3.	Mouse Citrate Plasma	-	0,099			
4.	Control: 1.+2. Antibody without antigen		0,096			
5.	Control: 1. Antibody without antigen	-	0,072			
6.	Control: 2. Antibody without antigen	-	0,085			
7.	Substrate (ABTS)	-	0,072			

Mouse plasma without the addition of human clotting factor IX showed an extinction of 0,099 at 405 nm background. When added human factor IX in a concentration of 2 mIU/ml (= 9 ng/ml human factor IX) the detection limit is reached. It can be deduced that the antihuman factor IX antibodies used in the ELISA are not cross-reactive with mouse coagulation factor IX.

Example 8: Cloning and Activity Testing of the Human Progesterone Receptor (hPR)

- 2. Activity Testing: The human progesterone receptor encoded in plasmid pTGhPR1 (s. Example 8.1 above) was tested for its phyiological activity. In a functional form and after activation with a progestin like R5020 the receptor should be able to induce the expression of luciferase from a Mouse Mammary Tumor Virus (MTV) promoter.
- To test this 293T cells were grown in phenol red-free DMEM 10 supplemented with 10% charcoal-filtrated fetal calf serum and with or without 10 nM of R5020 (NEN) in 6 well plates. Transfections were performed by the calcium phosphate method using 2 µg of a pSGhPR1 constructt and pMTV-luc (Hollenberg et al., 1985, Cell 55, p899-906) per well. One day after transfection the cells were washed in PBS 15 and the luciferase expression assayed with the Berthold luciferase kit according to the manufacturer's directions in а (Labsystems). The controls were as follows: R5020 was omitted (PR+MTV) and both plasmids alone were transfected with (PR+R5020, MTV+R5020) and without R5020 (PR, MTV). As positive control a 20 plasmid with a CMV-driven luciferase gene was transfected (pCMVluc).

As can be seen in Figure 19, there is a clear induction of luciferase expression when all the necessary elements are present, that is human progesterone receptor, progestin R5020 and the MTV-driven luciferase gene (PR+MTV+R5020). The error bars give the standard deviation of a threefold experiment, the readout is relative light units (RLU).

i 1.

Example 9: Oral Gene Transfer in in vivo Animal Experiment

<u>Purpose of experiment:</u> The object of this pilot study is to prove oral gene transfer in an in vivo animal experiment. Successful gene transfer is established by coagulation measurement: an additive effect of expressed human factor IX on the coagulation time of healthy murine whole blood is expected. The presence of expression of human factor IX in mouse blood is quantitated by ELISA.

Animals: The animals employed are 35 male C57BL/6J mice from Iffa Credo, France, with an initial age of 9 weeks and a weight of 23-33 g. The mice are kept in groups of 7 animals each in conventional test animal cages with wooden chips in the Institut für Experimentelle Onkologie und Therapieforschung der Technischen Universität München.

The animals are fed ad libitum with "Altrum Ratten und Mäuse Haltung" and are given tap water, also ad libitum.

The test animal cages are kept at an ambient temperature of 19-24°C and a humidity of 55 5%. The room is additionally provided with an automatic light supply which maintains a 12 hours rhythm.

The test animals are supervised by specialized staff.

Mixture of substances:

Group	Hormone	Hormone receptor	Plasmid	Aqua dest.	Route of administration
1	-	-	-	_	-
2	100 µl	-	10 µg	-	oral
3	-	-	10 µg	100 µl	oral
4	-	-	10 µg	50 µl	i.m.
5	100 μΙ	4.35 µg	10 µg	-	oral

Plasmid and hPR:

Theragene GmbH

Hormone:

Utrogest[®] by Dr. Kade/Besins Pharma GmbH,

Rigistr. 2, D-12277 Berlin

Aqua dest.:

5

10

15

20

25

30

Aqua ad injectabilia Delta-Pharma GmbH, 72793

Pfullingen

Esophageal sound: Vein catheter, diam. 0.5 x 0.9 mm.

Lot 7077 G2221, B. Braun Melsungen AG. Western

Germany

i.m. injection:

Micro-Fine 12.7 mm, Becton Dickinson GmbH,

Tullastr. 8-12, D-69126 Heidelberg

Course of experiment: The 35 mice were divided into 5 groups of 7 mice each. One group serves as a control, the second group was daily administered a total of 100 µl of hormone and plasmid via the gastrointestinal tract orally with an esophageal sound, the third group was daily administered a total of 100 µl of plasmid with aqua dest. orally with an esophageal sound, the fourth group was administered a total of 50 µl of plasmid with aqua dest. i.m. into the musculus quadriceps femoris, the fifth group was daily administered a total of 100 µl of hormone, hormone receptor and plasmid orally with an esophageal sound.

About 2-3 hours before the manipulation, the mice were prewarmed under a red light. Immediately before, during and after the manipulation, the mice were examined and supervised by a veterinarian.

Blood sampling from the mice was performed daily from the caudal artery of animals slightly sedated by inhalation anesthesia. For this purpuse the artery was punctured with a disposable injection cannula (0.90 x 40 mm). Whole blood welling out of the puncture site (5 µl of blood) was immediately collected with an Eppendorf pipette. Without further delay, the blood coagulation time in seconds was

15

20

25

30

determined using an Amelung-Koagulometer KC 4A by means of an aPTT assay (activated partial thromboplastin time). The blood coagulation analysis was always performed by the same person. Immediately after the blood sampling, the bleeding was stopped by compression.

Sedation of the mice was achieved by inhalation anesthesia (active substance: isoflurane: Forene, Abbott GmbH, 65205 Wiesbaden, Western Germany) in a whole body chamber.

The daily manipulation was performed through an overall period of 7 days. This was followed by a day (day 8 of experiment) without any manipulation, and at day 9 of experiment, again 5 µl of whole blood was withdrawn from the ventral caudal artery under anesthesia, and the coagulation time established as described above. Further, 0.5-0.75 ml of whole blood was collected intracardially using U-40 insulin syringes (Mikro-Fine 12.4 mm) filled with 50-75 µl of sodium citrate (3.1%), transferred into Eppendorf cuvettes, and about 100 µl of whole blood with citrate was reserved for PCR examination and stored in a cool environment. The remaining citrate blood was centrifuged for 10 min using a centrifuge 6000 rpm, 4°C, at 5000 rpm, and the plasma was recovered for the ELISA determination of the factor IX concentration.

Then, the animals were sacrificed using 0.5 ml Narkoren i.p. Immediately after the sacrificing, the animal bodies were dissected. The following organs were removed from the mice for an immunohistochemical examination: brain, spleen, liver, kidneys, testes, lungs, m. quadriceps femoris, heart, appendix; and frozen at -80°C.

<u>Deviation from the scheduled experimental course:</u> Due to the poor general condition of the mice in the course of the long-term administration series, the administration had to be interrupted at days

15

20

25

30

3 (except one mouse) and 5 for test group 2 (hormone and plasmid), at days 3 and 5 for group 5 (hormone, hormone receptor and plasmid), and two mice were additionally spared the administration of the reagents at days 2 and 7 of the experiment.

The poor general condition is accounted for by the hypnotic effect of the hormone progesterone. It causes the mice to sleep for about 24 hours without eating and drinking. This again has an adverse effect on the water balance of the mice, resulting in exsiccotic phenomena and apathic behavior. Therefore, the mice were prophylactically treated with a subcutaneous administration of 1 ml of 5% glucose solution (Delta Pharma GmbH, 72793 Pfullingen) and 1 ml of Ringer solution (Delta Pharma GmbH, 72793 Pfullingen) when the hormone was administered orally. Among the group which was orally administered hormone, hormone receptor and plasmid, two mice died at days 3 and 6, respectively; they were dissected.

Among the group which was orally administered hormone with plasmid, one mouse was found dead in its cage on day 8 of the experiment; it was also dissected.

The results are summarized in Figures 17 and 18. The statistical evaluations were performed according to the generalized linear model with repeated measurements (MANOVA with repeated measurements). In none of the test groups a non-linear course was observed. Therefore, the course was calculated by a simple representation of the linear increase or decrease, namely initial value minus final value per mouse. The particularly interesting difference between the control and the group "plasmid in the hormone with hormone receptor" (group 5) was examined using a T test for independent random samples.

Figure 17 shows the mean values of the calculated differences: In the control, for example, this difference was about 50 seconds. The vertical lines show plus and minus one standard deviation from these values. The T test is based both on the differences between the mean values and on the degree of overlapping which can be seen from these lines: The larger the overlapping, the less is the significance of the mean value differences. Thus, the groups "control" and "plasmid and water i.m." (groups 1 and 5, respectively) are distinguished in a purely numerical way in the mean value, but the degree of overlapping is so high that these groups are not significantly different.

The only significant difference was between group 1 and 5: The decrease of the latter is significantly higher than that of the control (T = -2.357; d.f. = 12; p < 0.05).

The following Tables contain the concluding statistics and the results of the statistical tests (T test) performed on the differences between the mean values obtained in the course of the test:

15

10

Group statistics

ADMI	N	N	mean value	standard deviation	standard error of the mean value
DIF	control	7	47.3857	58.9946	22.2978
	Hormone, hormone receptor and plasmid orally	7	114.7571	47.3300	17.8891

Test for independent random samples

1.			test for ariance							
		F	1 - 1 - 1 - 1 - 1 - 1		mean difference	standard error of difference	95% confidence interva of difference			
									lower	upper
DIF	variances are equal	0.026	0.874	-2.357	12	0.036	-67.3714	28.5869	-129.6570	-5.0858
	Variances are not equal			-2.357	11.461	0.037	-67.3714	28.5869	-129.9833	-4.7596

The human F IX was also detectable in the treated mice of the "hormone-hormone reception and plasmid orally group using an Elisa as described in Example 4.

Claims

- 1. Use of a nucleic acid construct comprising at least one hormone responsive element (HRE) and a transgene for preparing an agent for gene transfer.
- 2. The use of claim 1, wherein the at least one HRE is functionally linked to the transgene or not.
- 3. The use of claim 1 or 2, wherein the transgene is selected from the group consisting of genes encoding a blood clotting factor, hormone genes, hormone receptor genes, growth factors, enzyme genes, genes encoding cytokines or lymphokines, genes encoding inhibitor substances, genes encoding substances that function as drugs or vaccines, and antisense sequences.
 - 4. The use of claim 3, wherein the transgene is a gene encoding a blood clotting factor and the agent is suitable for treating hemophilia.
- 5. The of claim 4, wherein the human blood clotting factor is selected from the group consisting of factor VIII, factor IX, and von Willebrand Factor (vWF).
 - 6. The use of claims 1 to 5, wherein the nucleic acid construct comprises 1 to 20, preferably 3 to 10 HRE(s).
 - 7. The use of claim 1 to 5, wherein the at least one HRE is a steroid responsive element, preferably a progesterone responsive element (PRE).

- 8. The use of claim 5, wherein the HRE is a PRE and the blood clotting factor is factor IX, preferably the factor IX has a nucleotide sequence of 689 to 2071 of SEQ ID NO: 1.
- 9. The use of claim 5, wherein the HRE is a PRE and the blood clotting factor is factor VIII.
 - 10. The use of claim 7 to 9, wherein the PRE has the double stranded DNA sequence comprised of the DNA sequences of SEQ ID NOs: 3 and 4.
 - 11. The use of claims 1 to 10, wherein the construct further comprises functional DNA sequences selected from the group consisting of promoter sequences, enhancer sequences, silencer sequences, origin of replication sequences, integrational sequences, marker genes and switch sequences.
 - 12. The use of claim 11, wherein the construct further comprises a tissue-specific promoter, preferably an α -antitrypsin promoter.
 - 13. The use according to any one of claims 1 to 12, wherein the agent further comprises a hormone-hormone receptor complex, preferably a steroid-steroid receptor complex.
- 25 14. The use of claim 13, wherein the molar ratio of HRE within the nucleic acid construct to hormone receptor is from 1:1 to 1:10, preferably 1:2 to 1:5, and/or the molar ratio of hormone to hormone receptor is at least 1000:1, preferably at least 10000:1.
- 15. The use of claim 13 or 14, wherein the receptor is a progesterone

receptor and the steroid is progesterone or a progesterone derivative.

- 16. The use of claim 15, wherein the progesterone is natural micronized progesterone solubilized in a liphophilic matrix system and/or the progesterone receptor is hPR-A, hPR-B or comprises the nucleotide sequence of 557 to 933 SEQ ID NO: 9.
- 17. A nucleic acid construct comprising at least one HRE and a transgene, wherein one of said at least one HREs is not functionally linked to the transgene.
- 18. The nucleic acid construct of claim 17, which is as defined in claims 3 to 12.
- 19. A vector comprising the nucleic acid construct of claim 17 or 18.
 - 20. A transformed cell or transgenic organism comprising the nucleic acid construct as defined in claims 17 or 18 or the vector as defined in claim 19.

20

10

15

21. A composition of matter comprising a nucleic acid construct comprising at least one HRE and a transgene as defined in claim 17 or 18 and/or a vector as defined in claim 19, said at least one HRE being coupled to a hormone-hormone receptor complex.

- 22. The composition of matter of claim 21, wherein the hormone-hormone receptor complex is as defined in claims 13 to 16.
- 23. The composition of matter of claim 21, wherein the transgene is a gene encoding a blood clotting factor.

- 24. The composition of matter of claim 21 wherein the blood clotting factor is factor IX.
- 5 25. The composition of matter of claim 21 wherein the blood clotting factor is factor VIII.
 - 26. A method for preparing the composition of matter as defined in claim 21, which method comprises admixing the nucleic acid construct with the hormone receptor and the hormone.
 - 27. A pharmaceutical composition comprising the nucleic acid construct of claim 17 or 18, the vector of claim 19, and/or the composition of matter of claim 21 to 25.

25

- 28. The pharmaceutical composition of claim 27, which is suitable for gene transfer, preferably for treating hemophilia.
- 29. A method for gene transfer which comprises administering the agent as defined in claims 1 to 16, or the composition of matter as defined in claims 21 to 25 to an organism or to a cellular system.
 - 30. A method for delivering into an organism or into a cellular system a nucleic acid encoding a transgene to be expressed in the cells of the organism or the cells of the cellular system, which method comprises administering an agent as defined in claims 1 to 16 or composition of matter as defined in claims 21 to 25 to the organism or to the cellular system so that the hormone in the composition interacts with the cell membrane and therewith enhances diffusion and transport of the

20

nucleic acid that is coupled to the hormone-hormone receptor complex across the membrane and into the cell.

- 31. The method of claim 30, wherein a nucleic acid encoding human factor VIII or factor IX is delivered into the cell.
 - 32. A method of treating blood clotting disorders comprising administering a therapeutically effective amount of the composition of matter of claim 23 to an organism or to a cellular system.
 - 33. A method of treating hemophilia B, comprising administering a therapeutically effective amount of the composition of matter of claim 24 to an organism or to a cellular system.
- 34. A method of treating hemophilia A, comprising administering a therapeutically effective amount of the composition of matter of claim 25 to an organism or to a cellular system.
 - 35. Use of a steroid hormone for preparing an agent for gene transfer.
 - 36. The use of claim 35, wherein the steroid hormone is a natural micronized steroid hormone, preferably natural micronized progesterone.
- 25 37. The use of claim 36, wherein the natural micronized steroid hormone is solubilized in a lipophilic matrix system.
 - 38. A method for gene transfer which comprises administering a nucleic acid construct to an organism or to a cellular system, wherein

the nucleic acid construct contains a transgene and is encapsulated in a steroid hormone.




Fig. 2




Fig. 3




Fig. 4




Fig. 5




Fig. 6




Fig. 7





Fig. 8



9/22 Fig. 9

CGCGTTGACATTGATTATTGACTAGTTATTAATAGTAATCAATTACGGGGTCATTAGTTCATAGCCCATATATGGAGTTC TGTTCCCATAGTAACGCCAATAGGGACTTTCCATTGACGTCAATGGGTGGAGTATTTACGGTAAACTGCCCACTTGGCAG TACATCAAGTGTATCATATGCCAAGTACGCCCCCTATTGACGTCAATGACGGTAAATGGCCCGCCTGGCATTATGCCCAG TACATGACCTTATGGGACTTTCCTACTTGGCAGTACATCTACGTATTAGTCATCGCTATTACCATGGTGATGCGGTTTTG GCAGTACATCAATGGGCGTGGATAGCGGTTTGACTCACGGGGATTTCCAAGTCTCCACCCCATTGACGTCAATGGGAGTT TGTTTTGGCACCAAAATCAACGGGACTTTCCAAAATGTCGTAACAACTCCGCCCCATTGACGCAAATGGGCGGTAGGCGT GTACGGTGGGAGGTCTATATAAGCAGAGCTCTCTGGCTAACTAGAGAACCCACTGCTTACTGGCTTATCGAAATTAATAC GACTCACTATAGGGAGACCCAAGCTTGCATGCCAATTCCGCAAAGGTTATGCAGCGCGTGAACATGATCATGGCAGAATC ACCAGGCCTCATCACCATCTGCCTTTTAGGATATCTACTCAGTGCTGAATGTACAGTTTTTCTTGATCATGAAAACGCCA ATGGAAGAAAAGTGTAGTTTTGAAGAAGCACGAGAAGTTTTTGAAAAACACTGAAAGAACAACTGAATTTTGGAAGCAGTA TGTTGATGGAGATCAGTGTGAGTCCAATCCATGTTTAAATGGCGGCAGTTGCAAGGATGACATTAATTCCTATGAATGTT GGTGTCCCTTTGGATTTGAAGGAAAGAACTGTGAATTAGATGTAACATGTAACATTAAGAATGGCAGATGCGAGCAGTTT TGTAAAAATAGTGCTGATAACAAGGTGGTTTGCTCCTGTACTGAGGGATATCGACTTGCAGAAAACCAGAAGTCCTGTGA ACCAGCAGTGCCATTTCCATGTGGAAGAGTTTCTGTTTCACAAACTTCTAAGCTCACCCGTGCTGAGACTGTTTTTCCTG ATGTGGACTATGTAAATTCTACTGAAGCTGAAACCATTTTGGATAACATCACTCAAAGCACCCAATCATTTAATGACTTC ACTCGGGTTGTTGGTGGAGAAGATGCCAAACCAGGTCAATTCCCTTGGCAGGTTGTTTTGAATGGTAAAGTTGATGCATT CTGTGGAGGCTCTATCGTTAATGAAAAATGGATTGTAACTGCTGCCCACTGTGTTGAAACTGGTGTTAAAATTACAGTTG TCGCAGGTGAACATATATTGAGGAGACAGAACATACAGAGCAAAAGCGAAATGTGATTCGAATTATTCCTCACCACAAC TACAATGCAGCTATTAATAAGTACAACCATGACATTGCCCTTCTGGAACTGGACGAACCCTTAGTGCTAAACAGCTACGT TACACCTATTTGCATTGCTGACAAGGAATACACGAACATCTTCCTCAAATTTTGGATCTGGCTATGTAAGTGGCTGGGGAA GAGTCTTCCACAAAGGGAGATCAGCTTTAGTTCTTCAGTACCTTAGAGTTCCACTTGTTGACCGAGCCACATGTCTTCGA TCTACAAAGTTCACCATCTATAACAACATGTTCTGTGCTGGCTTCCATGAAGGAGGTAGAGATTCATGTCAAGGAGATAG TGGGGGACCCCATGTTACTGAAGTGGAAGGGACCAGTTTCTTAACTGGAATTATTAGCTGGGGTGAAGAGTGTGCAATGA AAGGCAAATATGGAATATATACCAAGGTATCCCGGTATGTCAACTGGATTAAGGAAAAAAACAAAGCTCACTTAATGGGAT CGGTCGAGCGGCCGCGACTCTACTAGAGGATCTTTGTGAAGGAACCTTACTTCTGTGGTGTGACATAATTGGACAAACTA CCTACAGAGATTTAAAGCTCTAAGGTAAATATAAAATTTTTAAGTGTATAATGTGTTAAACTACTGATTCTAATTGTTTG TGTATTTTAGATTCCAACCTATGGAACTGATGAATGGGAGCAGTGGTGGAATGCCTTTAATGAGGAAAACCTGTTTTGCT CAGAAGAAATGCCATCTAGTGATGATGAGGCTACTGCTGACTCTCAACATTCTACTCCTCCAAAAAAAGAAGAAAGGTA TGCTATTTACACCACAAAGGAAAAAGCTGCACTGCTATACAAGAAAATTATGGAAAAATATTCTGTAACCTTTATAAGTA GGCATAACAGTTATAATCATAACATACTGTTTTTTCTTACTCCACACAGGCATAGAGTGTCTGCTATTAATAACTATGCT CAAAAATTGTGTACCTTTAGCTTTTTAATTTGTAAAGGGGTTAATAAGGAATATTTGATGTATAGTGCCTTGACTAGAGA TCATAATCAGCCATACCACATTTGTAGAGGTTTTACTTGCTTTAAAAAACCTCCCACACCTCCCCCTGAACCTGAAACAT AAAATGAATGCAATTGTTGTTGTTAACTTGTTTATTGCAGCTTATAATGGTTACAAATAAAGCAATAGCATCACAAATTT CACAAATAAAGCATTTTTTTCACTGCATTCTAGTTGTGGTTTGTCCAAACTCATCAATGTATCTTATCATGTCTGGATCC CCGGGTACCCTCTAGAGCGAATTAATTCACTGGCCGTCGTTTTACAACGTCGTGACTGGGAAAACCCTGGCGTTACCCAA CTTAATCGCCTTGCAGCACATCCCCCTTTCGCCAGCTGGCGTAATAGCGAAGAGGCCCGCACCGATCGCCCTTCCCAACA GTTGCGCAGCCTGAATGGCGAATGGCGCCTGATGCGGTATTTTCTCCTTACGCATCTGTGCGGTATTTCACACCGCATAT TGACGGGCTTGTCTGCTCCCGGCATCCGCTTACAGACAAGCTGTGACCGTCTCCGGGAGCTGCATGTGTCAGAGGTTTTC ACCGTCATCACCGAAACGCGCGAGACGAAAGGGGGGGTACCAGCTTCGTAGCTAGAACATCATGTTCTGGGATATCAGCT TCGTAGCTAGAACATCATGTTCTGGTACCCCCCTCGTGATACGCCTATTTTTATAGGTTAATGTCATGATAATAATGGTT TCTTAGACGTCAGGTGGCACTTTTCGGGGAAATGTGCGCGGAACCCCTATTTGTTTATTTTTCTAAATACATTCAAATAT GTATCCGCTCATGAGACAATAACCCTGATAAATGCTTCAATAATATTGAAAAAGGAAGAGTATGAGTATTCAACATTTCC GTGTCGCCCTTATTCCCTTTTTTGCGGCATTTTGCCTTCCTGTTTTTGCTCACCCAGAAACGCTGGTGAAAGTAAAAGAT GCTGAAGATCAGTTGGGTGCACGAGTGGGTTACATCGAACTGGATCTCAACAGCGGTAAGATCCTTGAGAGTTTTCGCCC CGAAGAACGTTTTCCAATGATGAGCACTTTTAAAGTTCTGCTATGTGGCGCGGTATTATCCCGTATTGACGCCGGGCAAG AGCAACTCGGTCGCCGCATACACTATTCTCAGAATGACTTGGTTGAGTACTCACCAGTCACAGAAAAGCATCTTACGGAT CGGAGGACCGAAGGAGCTAACCGCTTTTTTGCACAACATGGGGGATCATGTAACTCGCCTTGATCGTTGGGAACCGGAGC TGAATGAAGCCATACCAAACGACGAGCGTGACACCACGATGCCTGTAGCAATGGCAACAACGTTGCGCAAACTATTAACT CTCGGCCCTTCCGGCTGGCTGTTTATTGCTGATAAATCTGGAGCCGGTGAGCGTGGGTCTCGCGGTATCATTGCAGCAC TGGGGCCAGATGGTAAGCCCTCCCGTATCGTAGTTATCTACACGACGGGGAGTCAGGCAACTATGGATGAACGAAATAGA CAGATCGCTGAGATAGGTGCCTCACTGATTAAGCATTGGTAACTGTCAGACCAAGTTTACTCATATATACTTTAGATTGA TTTAAAACTTCATTTTTAATTTAAAAGGATCTAGGTGAAGATCCTTTTTGATAATCTCATGACCAAAATCCCTTAACGTG GGTAACTGGCTTCAGCAGAGCGCAGATACCAAATACTGTTCTTCTAGTGTAGCCGTAGTTAGGCCACCACTTCAAGAACT GGGTTGGACTCAAGACGATAGTTACCGGATAAGGCGCAGCGGTCGGGCTGAACGGGGGGTTCGTGCACACAGCCCAGCTT GGAGCGAACGACCTACACCGAACTGAGATACCTACAGCGTGAGCTATGAGAAAGCGCCACGCTTCCCGAAGGGAGAAAGG

10/22

Fig. 9 (continued)


11/22 Fig. 10

Met 1	Gln	Arg	Val	Asn 5	Met	Ile	Met	Ala	Glu 10	Ser	Pro	Gly	Leu	Ile 15	Thr
Ile	Cys	Leu	Leu 20	Gly	Tyr	Leu	Leu	Ser 25	Ala	Glu	Cys	Thr	Val 30	Phe	Leu
Asp	His	Glu 35	Asn	Ala	Asn	Lys	Ile 40	Leu	Asn	Arg	Pro	Lys 45	Arg	Tyr	Asn
Ser	Gly 50	Lys	Leu	Glu	Glu	Phe 55	Val	Gln	Gly	Asn	Leu 60	Glu	Arg	Glu	Cys
Met 65	Glu	Glu	Lys	Cys	Ser 70	Phe	Glu	Glu	Ala	Arg 75	Glu	Val	Phe	Glu	Asn 80
Thr	Glu	Arg	Thr	Thr 85	Glu	Phe	Trp	Lys	Gln 90	Tyr	Val	Asp	Gly	Asp 95	Gln
Cys	Glu	Ser	Asn 100	Pro	Cys	Leu	Asn	Gly 105	Gly	Ser	Cys	Lys	Asp 110	Asp	Ile
Asn	Ser	Tyr 115	Glu	Cys	Trp	Cys	Pro 120	Phe	Gly	Phe	Glu	Gly 125	Lys	Asn	Cys
Glu	Leu 130		Val	Thr	Cys	Asn 135	Ile	Lys	Asn	Gly	Arg 140	Cys	Glu	Gln	Phe
Cys 145	Lys	Asņ	Ser	Ala	Asp 150	Asn	Lys	Val	Val	Cys 155	Ser	Cys	Thr	Glu	Gly 160
Tyr	Arg	Leu	Ala	Glu 165	Asn	Gln	Lys	Ser	Cys 170	Glu	Pro	Ala	Val	Pro 175	Phe
Pro	Cys	Gly	Arg 180	Val	Ser	Val	Ser	Gln 185	Thr	Ser	Lys	Leu	Thr 190	Arg	Ala
Glu	Thr	Val 195	Phe	Pro	Asp	Val	Asp 200	Tyr	Val	Asn	Ser	Thr 205	Glu	Ala	Glu
Thr	Ile 210	Leu	Asp	Asn	Ile	Thr 215	Gln	Ser	Thr		Ser 220	Phe	Asn	Asp	Phe
Thr 225	Arg	Val	Val	Gly	Gly 230	Glu	Asp	Ala	Lys	Pro 235	Gly	Gln	Phe	Pro	Trp 240
Gln	Val	Val	Leu	Asn 245	Gly	Lys	Val	Asp	Ala 250	Phe	Cys	Gly	Gly	Ser 255	Ile
Val	Asn	Glu	Lys 260	Trp	Ile	Val	Thr	Ala 265	Ala	His	Cys	Val	Glu 270	Thr	Gly
Val	Lys	Ile 275	Thr	Val	Val	Ala	Gly 280	Glu	His	Asn	Ile	Glu 285	Glu	Thr	Glu
His	Thr 290	Glu	Gln	Lys	Arg	Asn 295	Val	Ile	Arg	Ile	Ile 300	Pro	His	His	Asn

12/22 Fig. 10 (continued)

Tyr 305	Asn	Ala	Ala	Ile	Asn 310	Lys	Туr	Asn	His	Asp 315	Ile	Ala	Leu	Leu	Glu 320
Leu	Asp	Glu	Pro	Leu 325	Val	Leu	Asn	Ser	Tyr 330	Val	Thr	Pro	Ile	Cys 335	Ile
Ala	Asp	Lys	Glu 340	Tyr	Thr	Asn	Ile	Phe 345	Leu	Lys	Phe	Ġ1 y	Ser 350		Tyr
Val	Ser	Gly 355	Trp	Gly	Arg	Val	Phe 360	His	Lys	Gly	Arg	Ser 365	Ala	Leu	Val
Leu	Gln 370	Tyr	Leu	Arg	Val	Pro 375	Leu	Val	Asp	Arg	Ala 380	Thr	Cys	Leu	Arg
Ser 385	Thr	Lys	Phe	Thr	Ile 390	Tyr	Asn	Asn	Met	Phe 395	Cys	Ala	Gly	Phe	His 400
Glu	Gly	Gly	Arg	Asp 405	Ser	Cys	Gln	Gly	Asp 410	Ser	Gly	Gly	Pro	His 415	Val
Thr	Glu	Val	Glu 420	Gly	Thr	Ser	Phe	Leu 425	Thr	Gly	Ile	Ile	Ser 430	Trp	Gly
Glu	Glu	Cys 435	Ala	Met	Lys	Gly	Lys 440	Tyr	Gly	Ile	Tyr	Thr 445	Lys	Val	Ser
Arg	Tyr 450	Val	Asń	Trp	Ile	Lys 455	Glu	Lys	Thr	Lys	Leu 160	Thr.			

13/22 Fig. 11






Fig. 12a




Fig 12 b




Fig 12 c





Fig 12 d

16/22 Fig. **13**







Fig. 14



17/22

Fig. 15






Fig. 16









Fig. 18

Fig. 19



21/22

Fig. 20

```
ctgaccageg cegecetece eegeceeega eecaggaggt ggagateeet eeggteeage
    cacatteaac acceaettte tectecetet geceetatat tecegaaace ecetectet
121
    tecettttee etecteetg gagaeggggg aggagaaaag gggagteeag tegteatgae
    tgagctgaag gcaaagggtc cccgggctcc ccacgtggcg ggcggcccgc cctccccga
    ggtcggatcc ccactgctgt gtcgcccagc cgcaggtccg ttcccgggga gccagacctc
241
    ggacaccttg cctgaagttt cggccatacc tatctccctg gacgggctac tcttccctcg
301
361
    gccctgccag ggacaggacc cctccgacga aaagacgcag gaccagcagt cgctgtcgga
421
    cgtggagggc gcatattcca gagctgaagc tacaaggggt gctggaggca gcagttctag
481
    tcccccagaa aaggacagcg gactgctgga cagtgtcttg gacactctgt tggcgccctc
541
    aggtcccggg cagagccaac ccagcctcc cgcctgcgag gtcaccagct cttggtgcct
601
    gtttggcccc gaacttcccg aagatccacc ggctgccccc gccacccagc gggtgttgtc
    cccgctcatg agccggtccg ggtgcaaggt tggagacagc tccgggacgg cagctgccca
661
    taaagtgctg ccccggggcc tgtcaccagc ccggcagctg ctgctcccgg cctctgagag
721
781
    coctcactgg teeggggeee cagtgaagee gteteegeag geegetgegg tggaggttga
    ggaggaggat ggctctgagt ccgaggagtc tgcgggtccg cttctgaagg gcaaacctcg
841
    ggctctgggt ggcgcggcgg ctggaggagg agccgcggct gtcccgccgg gggcggcagc
901
961
    aggaggegte geeetggtee ceaaggaaga tteeegette teagegeeea gggtegeeet
1021 ggtggagcag gacgcgccga tggcgcccgg gcgctccccg ctggccacca cggtgatgga
1081 tttcatccac gtgcctatcc tgcctctcaa tcacgcctta ttggcagccc gcactcggca
1141 gctgctggaa gacgaaagtt acgacggcgg ggccggggct gccagcgcct ttgccccgcc
1201 gcggagttca ccctgtgcct cgtccacccc ggtcgctgta ggcgacttcc ccgactgcgc
1261 gtacccgccc gacgccgagc ccaaggacga cgcgtaccct ctctatagcg acttccagcc
1321 gcccgctcta aagataaagg aggaggagga aggcgcggag gcctccgcqc qctccccqcq
1381 ttcctacctt gtggccggtg ccaaccccgc agccttcccg gatttcccgt tggggccacc
1441 gcccccgctg ccgccgcgag cgaccccatc cagacccggg gaagcggcgg tgacggccgc
1501 acceptcagt gentcagtet egtetgegte etcetegggg tegaceetgg agtgeateet
1561 gtacaaageg gagggegege egeeceagea gggeeegtte gegeegeege eetgeaagge
1621 gccgggcgcg agcggctgcc tgctcccgcg ggacggcctg ccctccacct ccgcctctgc
1681 egeogeogeo ggggeggee cegegeteta ecetgeacte ggeeteaaeg ggeteeegea
1741 gctcggctac caggccgccg tgctcaagga gggcctgccg caggtctacc cgccctatct
1801 caactacctg aggeoggatt cagaagccag ccagagccca caatacagct tcgagtcatt
1861 acctcagaag atttgtttaa tctgtgggga tgaagcatca ggctgtcatt atggtgtcct
1921 tacctgtggg agctgtaagg tcttctttaa gagggcaatg gaagggcagc acaactactt
1981 atgtgctgga agaaatgact gcatcgttga taaaatccgc agaaaaaact gcccagcatg
2041 tegeettaga aagtgetgte aggetggeat ggteettgga ggtegaaaat ttaaaaagtt
2101 caataaagtc agagttgtga gagcactgga tgctgttgct ctcccacagc cattgggcgt
2161 tocaaatgaa agccaagcco taagccagag attcactttt toaccaggto aagacataca
2221 gttgattcca ccactgatca acctgttaat gagcattgaa ccagatgtga tctatgcagq
2281 acatgacaac acaaaacctg acacctccag ttctttgctg acaagtctta atcaactagg
2341 cgagaggcaa cttctttcag tagtcaagtg gtctaaatca ttgccaggtt ttcgaaactt
2401 acatattgat gaccagataa ctctcattca gtattcttgg atgagcttaa tggtgtttgg
2461 totaggatgg agatoctaca aacatgtoag tgggcagatg otgtattttg cacotgatot
2521 aatactaaat gaacagegga tgaaagaate atcattetat teattatgee ttaccatgtg
2581 gcagatecca caggagtttg teaagettea agttagecaa gaagagttee tetgtatgaa
2641 agtattgtta cttcttaata caattccttt ggaagggcta cgaagtcaaa cccagtttga
2701 ggagatgagg tcaagctaca ttagagagct catcaaggca attggtttga ggcaaaaagg
2761 agttgtgtcg agctcacagc gtttctatca acttacaaaa cttcttgata acttgcatga
2821 tettgicaaa caacticate tgtactgett gaatacattt atccagtece gggeactgag
2881 tgttgaattt ccagaaatga tgtctgaagt tattgctgca caattaccca agatattggc
2941 agggatggtg aaaccccttc tctttcataa
```

Fig. 21

```
SEQUENCE LISTING
    <110> Theragene Biomedical Laboratories GmbH
    <120> Hormone-Hormone Receptor Complexes and Nucleic Acid
          Constructs and Their Use in Gene Therapy
    <130> 000065wo/JH/ml
10
    <140>
    <141>
    <160> 19
15
    <170> PatentIn Ver. 2.1
    <210> 1
     <211> 5753
20
     <212> DNA
    <213> Artificial Sequence
    <223> Description of Artificial Sequence: vector pTGFG36
25
    <220>
     <221> CDS
     <222> (689)..(2071)
30
    cgcgttgaca ttgattattg actagttatt aatagtaatc aattacgggg tcattagttc 60
    atageceata tatggagtte egegttaeat aacttaeggt aaatggeeeg eetggetgae 120
35
    cgcccaacga ccccgccca ttgacgtcaa taatgacgta tgttcccata gtaacgccaa 180
    tagggacttt ccattgacgt caatgggtgg agtatttacg gtaaactgcc cacttggcag 240
    tacatcaagt gtatcatatg ccaagtacgc cccctattga cgtcaatgac ggtaaatggc 300
40
    ccgcctggca ttatgcccag tacatgacct tatgggactt tcctacttgg cagtacatct 360
    acgtattagt catcgctatt accatggtga tgcggttttg gcagtacatc aatgggcgtg 420
45
    gatageggtt tgacteaegg ggattteeaa gteteeaece cattgaegte aatgggagtt 480
    tgttttggca ccaaaatcaa cgggactttc caaaatgtcg taacaactcc gccccattga 540
    cgcaaatggg cggtaggcgt gtacggtggg aggtctatat aagcagagct ctctggctaa 600
50
    ctagagaacc cactgcttac tggcttatcg aaattaatac gactcactat agggagaccc 660
    aagcttgcat gccaattccg caaaggtt atg cag cgc gtg aac atg atc atg
                                                                        712
                                    Met Gln Arg Val Asn Met Ile Met
55
    gca gaa tca cca ggc ctc atc acc atc tgc ctt tta gga tat cta ctc
                                                                        760
    Ala Glu Ser Pro Gly Leu Ile Thr Ile Cys Leu Leu Gly Tyr Leu Leu
          10
                              15
60
```

	23		aca gtt Thr Val			35	Ala Asn	Lys Ile	
			45		50	тур пец	Giu Giu	Phe Val	856
10	caa ggg Gln Gly	g aac ctt / Asn Leu 60	gag aga Glu Arg	gaa tgt a Glu Cys N	atg gaa Met Glu 65	gaa aag Glu Lys	tgt agt 1 Cys Ser 1 70		904
· 15	gaa gca Glu Ala	cga gaa Arg Glu 75	gtt ttt (Val Phe (gaa aac a Glu Asn T 80	ict gaa hr Glu	aga aca a Arg Thr !		tt tgg he Trp	952
20	aag cag Lys Gln 90	tat gtt Tyr Val	gat gga g Asp Gly A	gat cag t Asp Gln C 95	gt gag ys Glu	tcc aat o Ser Asn I	-	ta aat eu Asn	1000
	ggc ggc Gly Gly 105	agt tgc Ser Cys	aag gat g Lys Asp A 110	ac att a sp Ile A			gt tgg t ys Trp C	ys Pro	1048
25	ttt gga Phe Gly	ttt gaa Phe Glu	gga aag a Gly Lys A 125	ac tgt ga sn Cys G			nr Cys As	sn Ile	1096
30	aag aat Lys Asn	ggc aga Gly Arg 140	tgc gag ca Cys Glu G	ag ttt to ln Phe Cy 14	jt aaa a 's Lys A	aat agt g Asn Ser A	ct gat aa la Asp As	35 ac aag an Lys	1144
35		155	tgt act ga Cys Thr Gl	160 1	9 15	icu Ara G	lu Asn Gl 55	n Lys	1192
40	170		gca gtg co Ala Val Pr 17	5	- J - C	ga aga gt ly Arg Va 180	t tct gt 1 Ser Va	l Ser	1240
	185		etc acc cg Leu Thr Ar 190		19	95	o Asp Va	l Asp	1288
45	.:	2	ct gaa gc hr Glu Ala 05		210	ASP AS	o TTe Thi	caa Gln	1336
50	agc acc c Ser Thr G	aa tca t In Ser P 220	tt aat gad he Asn Asp	c ttc act Phe Thr 225	- 5 · u	t gtt gg			1384
55	gcc aaa c Ala Lys P 2	ca ggt ca ro Gly G 35	aa ttc cct ln Phe Pro	tgg cag Trp Gln 240	gtt gt Val Va	t ttg aat 1 Leu Asr 245	ggt aaa Gly Lys	gtt Val	1432
60	gat gca t Asp Ala Pl 250	tc tgt go he Cys Gl	ga ggc tct ly Gly Ser 255	atc gtt Ile Val	aat ga Asn Gl	_		act Thr	1480

												aca Thr					1528
. 5	gaa Glu	cat His	aat Asn	att Ile	gag Glu 285	gag Glu	aca Thr	gaa Glu	cat His	aca Thr 290	gag Glu	caa Gln	aag Lys	cga Arg	aat Asn 295	gtg Val	1576
10												gct Ala					1624
15												ccc Pro					1672
20								Ile				gaa Glu 340					1720
20												tgg Trp					1768
25												ctt Leu				Leu	1816
30	gtt Val											ttc Phe					1864
35												aga Arg					1912
40												gaa Glu 420					1960
40												gca Ala					2008
45	tat Tyr	gga Gly	ata Ile	tat Tyr	acc Thr 445	aag Lys	gta Val	tcc Ser	cgg Arg	tat Tyr 450	Val	aac Asn	tgg Trp	att Ile	aag Lys 455	gaa Glu	2056
50			aag Lys		Thr	taa	tggg	atc	ggtc	gagc	gg c	cgcg	actc	t ac	taga	ggat	2111
	ctt	tgtg	aag	gaac	ctta	ct t	ctgt	ggtg	t ga	cata	attg	gac	aaac	tac	ctac	agagat	2171
55	tta	aagc	tct	aagg	taaa	ta t	aaaa	tttt	t aa	gtgt	ataa	tgt	gtta	aac	tact	gattct	2231
	aat	tgtt	tgt	gtat	ttta	ga t	tcca	acct	a tg	gaac	tgat	gaa	tggg	agc	agtg	gtggaa	2291
60	tgc	cttt	aat	gagg	aaaa	cc t	gttt	tgct	c ag	aaga	aatg	cca	tcta	gtg	atga	tgaggc	2351
	tac	tgct	gac	tctc	aaca	tt c	tact	cctc	c aa	aaaa	gaag	aga	aagg	tag	aaga	ccccaa	2411

	*ggactttcct	tcagaattgc	taagttttt	gagtcatgct	gtgtttagta	atagaactct	2471
	tgcttgcttt	gctatttaca	ccacaaagga	aaaagctgca	ctgctataca	agaaaattat	2531
5	ggaaaaatat	tctgtaacct	ttataagtag	gcataacagt	tataatcata	acatactgtt	2591
	ttttcttact	ccacacaggc	atagagtgtc	tgctattaat	aactatgctc	aaaaattgtg	2651
0	tacctttagc	tttttaattt	gtaaaggggt	taataaggaa	tatttgatgt	atagtgcctt	2711
	gactagagat	cataatcagc	cataccacat	ttgtagaggt	tttacttgct	ttaaaaaacc	2771
	·tcccacacct	cccctgaac	ctgaaacata	aaatgaatgc	aattgttgtt	gttaacttgt	2831
5·	:ttattgcagc	ttataatggt	tacaaataaa	gcaatagcat	cacaaatttc	acaaataaag	2891
	cattttttc	actgcattct	agttgtggtt	tgtccaaact	catcaatgta	tcttatcatg	2951
20	tctggatccc	cgggtaccct	ctagagcgaa	ttaattcact	ggccgtcgtt	ttacaacgtc	3011
.0	gtgactggga	aaaccctggc	gttacccaac	ttaatcgcct	tgcagcacat	cccctttcg	3071
	ccagctggcg	taatagcgaa	gaggcccgca	ccgatcgccc	ttcccaacag	ttgcgcagcc	3131
25	tgaatggcga	atggcgcctg	atgcggtatt	ttctccttac	gcatctgtgc	ggtatttcac	3191
	accgcatatg	gtgcactctc	agtacaatct	gctctgatgc	cgcatagtta	agccagcccc	3251
80	gacacccgcc	aacacccgct	gacgcgccct	gacgggcttg	tctgctcccg	gcatccgctt	3311
	acagacaagc	tgtgaccgtc	tccgggagct	gcatgtgtca	gaggttttca	ccgtcatcac	3371
	cgaaacgcgc	gagacgaaag	ggggggtacc	agcttcgtag	ctagaacatc	atgttctggg	3431
15	atatcagctt	cgtagctaga	acatcatgtt	ctggtacccc	cctcgtgata	cgcctatttt	3491
	tataggttaa	tgtcatgata	ataatggttt	cttagacgtc	aggtggcact	tttcggggaa	3551
10	atgtgcgcgg	aacccctatt	tgtttatttt	tctaaataca	ttcaaatatg	tatccgctca	3611
	tgagacaata	accctgataa	atgcttcaat	aatattgaaa	aaggaagagt	atgagtattc	3671
	aacatttccg	tgtcgccctt	attccctttt	ttgcggcatt	ttgccttcct	gtttttgctc	3731
5	acccagaaac	gctggtgaaa	gtaaaagatg	ctgaagatca	gttgggtgca	cgagtgggtt	3791
	acatcgaact	ggatctcaac	agcggtaaga	tccttgagag	ttttcgcccc	gaagaacgtt	3851
50	ttccaatgat	gagcactttt	aaagttctgc	tatgtggcgc	ggtattatcc	cgtattgacg	3913
-	ccgggcaaga	gcaactcggt	cgccgcatac	actattctca	gaatgacttg	gttgagtact	3971
	caccagtcac	agaaaagcat	cttacggatg	gcatgacagt	aagagaatta	tgcagtgctg	4031
55	ccataaccat	gagtgataac	actgcggcca	acttacttct	gacaacgatc	ggaggaccga	4091
	aggagctaac	cgcttttttg	cacaacatgg	gggatcatgt	aactcgcctt	gatcgttggg	415
50	aaccggagct	gaatgaagcc	ataccaaacg	acgagcgtga	caccacgatg	cctgtagcaa	421
-	tggcaacaac	gttgcgcaaa	ctattaactg	gcgaactact	tactctagct	tcccggcaac	427

	aattaataga	ctggatggag	gcggataaag	ttgcaggaco	acttctgcgc	teggeeette	4331
	cggctggctg	gtttattgct	gataaatctg	gagccggtga	gcgtgggtct	cgcggtatca	4391
. 5	ttgcagcact	ggggccagat	ggtaagccct	cccgtatcgt	agttatctac	acgacgggga	4451
	gtcaggcaac	tatggatgaa	cgaaatagac	agatcgctga	gataggtgcc	tcactgatta	4511
10			caagtttact				
	atttttaatt	taaaaggatc	taggtgaaga	tcctttttga	taatctcatg	accaaaatcc	4631
	cttaacgtga	gttttcgttc	cactgagegt	cagaccccgt	agaaaagatc	aaaggatctt	4691
15	cttgagatcc	tttttttctg	cgcgtaatct	gctgcttgca	aacaaaaaa	ccaccgctac	4751
	cagcggtggt	ttgtttgccg	gatcaagagc	taccaactct	ttttccgaag	gtaactggct	4811
20			aatactgttc				
	tcaagaactc	tgtagcaccg	cctacatacc	tegetetget	aatcctgtta	ccagtggctg	4931
	ctgccagtgg	cgataagtcg	tgtcttaccg	ggttggactc	aagacgatag	ttaccggata	4991
2 5			acggggggtt		•		
	cctacaccga	actgagatac	ctacagcgtg	agctatgaga	aagcgccacg	cttcccgaag	5111
30	ggagaaaggc	ggacaggtat	ccggtaagcg	gcagggtcgg	aacaggagag	cgcacgaggg	5171
	agcttccagg	gggaaacgcc	tggtatcttt	atagtcctgt	cgggtttcgc	cacctctgac_	5231
	ttgagcgtcg	atttttgtga	tgctcgtcag	gggggcggag	cctatggaaa	aacgccagca	5291
35	acgcggcctt	tttacggttc	ctggcctttt	gctggccttt	Pgctcacatg	ttctttcctg	5351
	cgttatcccc	tgattctgtg	gataaccgta	ttaccgcctt	tgagtgagct	gataccgctc	5411
40	gccgcagccg	aacgaccgag	cgcagcgagt	cagtgagcga	ggaagcggaa	gagcgcccaa	5471
	tacgcaaacc	gcctctcccc	gcgcgttggc	cgattcatta	atgcagctgg	cacgacaggt	5531
	ttcccgactg	gaaagcgggc	agtgagcgca	acgcaattaa	tgtgagttag	ctcactcatt	5591
45	aggcacccca	ggctttacac	tttatgcttc	cggctcgtat	gttgtgtgga	attgtgagcg	5651
			aacagctatg				
50	tagagctcta	gagctctaga	gagcttgcat	gcctgcaggt	cg		5753
55	<223> Descr	ficial Seque ription of A	ence Artificial S	equence: ve	ctor pTGFG3	6	
6 0	<400> 2 Met Gln Arg	y Val Asn Me 5	t Ile Met A	la Glu Ser 10	Pro Gly Leu	Ile Thr	

	Ile	e Cys	s Leu	Let 20	ı Gly	' Tyr	Leu	ı Lev	Ser 25	Ala	a Glu	Cys	Thr	: Val		Leu
5	Asp	His	35 35	Asr	Ala	Asn	Lys	3 Ile 40	Leu	Asn	Arg	Pro	Lys 45	Arg	Tyr	Asn
	Ser	Gl ₃ 50	Lys	Leu	Glu	Glu	Phe 55	val	Gln	Gly	/ Asn	Leu 60	Glu	Arg	Glu	Cys
10	Met 65	Glu	a Glu	Lys	Cys	Ser 70	Phe	Glu	Glu	Ala	Arg 75	Glu	Val	Phe	Glu	Asn 80
15	Thr	Glu	Arg	Thr	Thr 85	Glu	Phe	Trp	Lys	Gln 90	Tyr	Val	Asp	Gly	Asp 95	Gln
	Cys	Glu	Ser	Asn 100	Pro	Cys	Leu	Asn	Gly 105	Gly	Ser	Cys	Lys	Asp 110	Asp	Ile
20	Asn	Ser	Tyr 115	Glu	Cys	Trp	Cys	Pro 120	Phe	Gly	Phe	Glu	Gly 125	Lys	Asn	Cys
•	Glu	Leu 130	Asp	Val	Thr	Cys	Asn 135	Ile	Lys	Asn	Gly	Arg 140	Cys	Glu	Gln	Phe
25	Cys 145	Lys	Asn	Ser	Ala	Asp 150	Asn	Lys	Val	Val	Cys 155	Ser	Cys	Thr	Glu	Gly 160
30	Tyr	Arg	Leu	Ala	Glu 165	Asn	Gln	Lys	Ser	Cys 170	Glu	Pro	Ala	Val	Pro 175	Phe
	Pro	Cys	Gly	Arg 180	Val	Ser	Val	Ser	Gln 185	Thr	Ser	Lys	Leu	Thr 190	Arg	Ala
35	Glu	Thr	Val 195	Phe	Pro	Asp	Val	Asp -200	Tyr	Val	Asn	Ser	Thr 205	Glu	Ala	Glu
	Thr	Ile 210	Leu	Asp	Asn	Ile	Thr 215	Gln	Ser	Thr	Gln	Ser 220	Phe	Asn	Asp	Phe
40	Thr 225	Arg	Val	Val	Gly	Gly 230	Glu	Asp	Ala	Lys	Pro 235	Gly	Gln	Phe	Pro	Trp 240
45	Gln	Val	Val	Leu	Asn 245	Gly	Lys	Val	Asp	Ala 250	Phe	Cys	Gly	Gly	Ser 255	Ile
	Val	Asn	Glu	Lys 260	Trp	Ile	Val	Thr	Ala 265	Ala	His	Cys	Val	Glu 270	Thr	Gly
50	Val	Lys	Ile 275	Thr	Val	Val	Ala	Gly 280	Glu	His	Asn	Ile	Glu 285	Glu	Thr	Glu
	His	Thr 290	Glu	Gln	Lys	Arg	Asn 295	Val	Ile	Arg	Ile	Ile 300	Pro	His	His	Asn
5 5	Tyr 305	Asn	Ala	Ala	Ile	Asn 310	Lys	Tyr	Asn	His	Asp 315	Ile	Ala	Leu		Glu 320
6 0	Leu	Asp	Glu	Pro	Leu 325	Val	Leu	Asn	Ser	Tyr 330	Val	Thr	Pro		Cys 335	Ile
	Ala	Asp	Lys	Glu 340	Tyr	Thr	Asn	Ile	Phe 345	Leu	Lys	Phe		Ser 350	Gly	Tyr

	Val	Ser	Gly 355	Trp	Gly [.]	Arg	Val	Phe 360	His	Lys	Gly	Arg	Ser 365	Ala	Leu	Val	
5	Leu	Gln 370	Tyr	Leu	Arg	Val	Pro 375	Leu	Val	Asp	Arg	Ala 380	Thr	Cys	Leu	Arg	
10	Ser 385	Thr	Lys	Phe	Thr	Ile 390	Tyr	Asn	Asn	Met	Phe 395	Cys	Ala	Gly	Phe	His 400	
	Glú	Gly	Gly	Arg	Asp 405	Ser	Cys	Gln	Gly	Asp 410	Ser	Gly	Gly	Pro	His 415	Val	
15	Thr	Glu	Val	Glu 420	Gly	Thr	Ser	Phe	Leu 425	Thr	Gly	Ile	Ile	Ser 430	Trp	Gly	
	Glu	Glu	Cys 435	Ala	Met	Lys	Gly	Lys 440	Tyr	Gly	Ile	Tyr	Thr 445	Lys	Val	Ser	
20	Arg	Tyr 450	Val	Asn	Trp	Ile	Lys 455	Glu	Lys	Thr	Lys	Leu 460	Thr				
25	<212 <212	0> 3 1> 78 2> Di 3> Ho	ΝA	sapie	ens												. ,
30	gggg	0> 3 gtaco atgtt	cag	ette	gtag	ct ac	gaaca	atcai	gt1	ctg	ggat	atca	agcti	tcg 1	tagci	tagaac	60 78
35	<212	0> 4 1> 78 2> DN 3> Ho	AV	sapie	ens												
40	ggg	0> 4 gtaco cgaaq				gt to	ctago	ctac	g aaq	gctga	atat	ccca	agaa	cat o	gatgi	ttctag	60 78
45	<212	0> 5 1> 19 2> DI 3> Ho	AV	sapi	ens			•	·. -	.÷	· .						
50	<400 agc	0> 5 ttgad	cct (cgag	caag	С											19
55	<21:	0> 6 1> 1: 2> DI 3> He	NA	sapi	ens												
6 0		0> 6 cgct	tgc	tcga	ggtc	a											19

```
<210> 7
      <211> 43
      <212> DNA
      <213> Homo sapiens
      <400> 7
      ggaattccgc aaaggttatg cagcgcgtga acatgatcat ggc
                                                                           43
 10
      <210> 8
      <211> 39
      <212> DNA
      <213> Homo sapiens
 15
     cgcggatcca ttaagtgagc tttgtttttt ccttaatcc
                                                                           39
     <210> 9
 20
     <211> 26
     <212> DNA
     <213> Homo sapiens
     <400> 9
     cgaggatcca gtcgtcatga ctgagc
                                                                           26
     <210> 10
     <211> 41
     <212> DNA
30
     <213> Homo sapiens
     <400> 10
     gcagaattca ttataaaaac tcaagacctc ataatcctga c
35
                                                                          41
     <210> 11
     <211> 20
     <212> DNA
40
     <213> Homo sapiens
     <400> 11
     ctcctcgggg tcgaccctgg
                                                                          20
45
     <210> 12
     <211> 20
     <212> DNA
     <213> Homo sapiens
50
     <400> 12
    ccagggtcga ccccgaggag
                                                                          20
55
    <210> 13
    <211> 5905
    <212> DNA
    <213> Artificial Sequence
60
    <220>
    <223> Description of Artificial Sequence: vector pTGFG53
```

```
<400> 13
     cgcgttgaca ttgattattg actagttatt aatagtaatc aattacgggg tcattagttc 60
     atageceata tatggagtte egegttaeat aacttaeggt aaatggeeeg eetggetgae 120
     cgcccaacga cccccgccca ttgacgtcaa taatgacgta tgttcccata gtaacgccaa 180
     tagggacttt ccattgacgt caatgggtgg agtatttacg gtaaactgcc cacttggcag 240
     tacatcaagt gtatcatatg ccaagtacgc cccctattga cgtcaatgac ggtaaatggc 300
     ccgcctggca ttatgcccag tacatgacct tatgggactt tcctacttgg cagtacatct 360
     acgtattagt catcgctatt accatggtga tgcggttttg gcagtacatc aatgggcgtg 420
     gatageggtt tgacteaegg ggattteeaa gteteeaece cattgaegte aatgggagtt 480
10
     tgttttggca ccaaaatcaa cgggactttc caaaatgtcg taacaactcc gccccattga 540
     cgcaaatggg cggtaggcgt gtacggtggg aggtctatat aagcagagct ctctggctaa 600
     ctagagaacc cactgettae tggettateg aaattaatac gacteactat agggagaecc 660
     aagettgeat gecaatteeg caaaggttat geagegegtg aacatgatea tggeagaate 720
     accaggeete atcaccatet geettttagg atatetaete agtgetgaat gtacagtttt 780
15
     tcttgatcat gaaaacgcca acaaaattct gaatcggcca aagaggtata attcaggtaa 840
     attggaagag tttgttcaag ggaaccttga gagagaatgt atggaagaaa agtgtagttt 900
     tgaagaagca cgagaagttt ttgaaaacac tgaaagaaca actgaatttt ggaagcagta 960
     tgttgatgga gatcagtgtg agtccaatcc atgtttaaat ggcggcagtt gcaaggatga 1020
     cattaattcc tatgaatgtt ggtgtccctt tggatttgaa ggaaagaact gtgaattaga 1080
20
     tgtaacatgt aacattaaga atggcagatg cgagcagttt tgtaaaaata gtgctgataa 1140
     caaggtggtt tgctcctgta ctgagggata tcgacttgca gaaaaccaga agtcctgtga 1200
     accagcagtg ccatttccat gtggaagagt ttctgtttca caaacttcta agctcacccg 1260
     tgctgagact gtttttcctg atgtggacta tgtaaattct actgaagctg aaaccatttt 1320
     ggataacatc actcaaagca cccaatcatt taatgacttc actcgggttg ttggtggaga 1380
25
     agatgecaaa ecaggteaat teeettggea ggttgttttg aatggtaaag ttgatgeatt 1440
     ctgtggaggc tctatcgtta atgaaaaatg gattgtaact gctgcccact gtgttgaaac 1500
     tggtgttaaa attacagttg tcgcaggtga acataatatt gaggagacag aacatacaga 1560
     gcaaaagcga aatgtgattc gaattattcc tcaccacaac tacaatgcag ctattaataa 1620
     gtacaaccat gacattgccc ttctggaact ggacgaaccc ttagtgctaa acagctacgt 1680
30
     tacacctatt tgcattgctg acaaggaata cacgaacatc ttcctcaaat ttggatctgg 1740
     ctatgtaagt ggctggggaa gagtcttcca caaagggaga tcagctttag ttcttcagta 1800
     ccttagagtt ccacttgttg accgagccac atgtcttcga tctacaaagt tcaccatcta 1860
     taacaacatg ttetgtgetg gettecatga aggaggtaga gatteatgte aaggagatag 1920
     tgggggaccc catgttactg aagtggaagg gaccagtttc ttaactggaa ttattagctg 1980
35
     gggtgaagag tgtgcaatga aaggcaaata tggaatatat accaaggtat cccggtatgt 2040
     caactggatt aaggaaaaaa caaagctcac ttaatgggat cggtcgagcg gccgcgactc 2100
     tactagagga tetttgtgaa ggaacettae ttetgtggtg tgacataatt ggacaaacta 2160
     cctacagaga tttaaagctc taaggtaaat ataaaatttt taagtgtata atgtgttaaa 2220
     ctactgattc taattgtttg tgtattttag attccaacct atggaactga tgaatgggag 2280
40
     cagtggtgga atgcctttaa tgaggaaaac ctgttttgct cagaagaaat gccatctagt 2340
     gatgatgagg ctactgctga ctctcaacat tctactcctc caaaaaagaa gagaaaggta 2400
     gaagacccca aggactttcc ttcagaattg ctaagttttt tgagtcatgc tgtgtttagt 2460
     aatagaacte tigettgett tgetatttae accaeaagg aaaaagetge actgetatae 2520
     aagaaaatta tggaaaaata ttctgtaacc tttataagta ggcataacag ttataatcat 2580
45
     aacatactgt tttttcttac tccacacagg catagagtgt ctgctattaa taactatgct 2640
     caaaaattgt gtacctttag ctttttaatt tgtaaagggg ttaataagga atatttgatg 2700
     tatagtgcct tgactagaga tcataatcag ccataccaca tttgtagagg ttttacttgc 2760
     tttaaaaaac ctcccacacc tccccctgaa cctgaaacat aaaatgaatg caattgttgt 2820
     tgttaacttg tttattgcag cttataatgg ttacaaataa agcaatagca tcacaaattt 2880
50
     cacaaataaa gcattttttt cactgcattc tagttgtggt ttgtccaaac tcatcaatgt 2940
     atcttatcat gtctggatcc ccggggggta ccagcttcgt agctagaaca tcatgttctg 3000
     ggatatcagc ttcgtagcta gaacatcatg ttctggtacc cccgctctag agcgaattaa 3060
     ttcactggcc gtcgttttac aacgtcgtga ctgggaaaac cctggcgtta cccaacttaa 3120
     tegeettgea geacateece etttegeeag etggegtaat agegaagagg eeegcacega 3180
55
     tegecettee caacagttge geageetgaa tggcgaatgg egeetgatge ggtatttet
     cettaegeat etgtgeggta ttteacaceg catatggtge acteteagta caatetgete 3300
     tgatgccgca tagttaagcc agccccgaca cccgccaaca cccgctgacg cgccctgacg 3360
     ggcttgtctg ctcccggcat ccgcttacag acaagctgtg accgtctccg ggagctgcat 3420
     gtgtcagagg ttttcaccgt catcaccgaa acgcgcgaga cgaaagggcg gggtaccaga 3480
60
     acatgatgtt ctagctacga agetgatate ccagaacatg atgttetage tacgaagetg 3540 gtacccegge etegtgatae geetatttt ataggttaat gteatgataa taatggttte 3600
     ttagacgtca ggtggcactt ttcggggaaa tgtgcgcgga acccctattt gtttattttt 3660
```

```
ctaaatacat tcaaatatgt atccgctcat gagacaataa ccctgataaa tgcttcaata 3720
     atattgaaaa aggaagagta tgagtattca acatttccgt gtcgccctta ttcccttttt 3780
     tgcggcattt tgccttcctg tttttgctca cccagaaacg ctggtgaaag taaaagatgc 3840
     tgaagatcag ttgggtgcac gagtgggtta catcgaactg gatctcaaca gcggtaagat 3900
     cettgagagt tttegeceeg aagaaegttt tecaatgatg ageaetttta aagttetget 3960
 5
     atgtggcgcg gtattatccc gtattgacgc cgggcaagag caactcggtc gccgcataca 4020
     ctatteteag aatgaettgg ttgagtaete accagteaea gaaaageate ttaeggatgg 4080
     catgacagta agagaattat gcagtgctgc cataaccatg agtgataaca ctgcggccaa 4140
     cttacttctg acaacgateg gaggacegaa ggagetaacc gcttttttgc acaacatggg 4200
     ggatcatgta actegeettg ategttggga aceggagetg aatgaageea taccaaaega 4260
10
     cgagcgtgac accacgatgc ctgtagcaat ggcaacaacg ttgcgcaaac tattaactgg 4320
     cgaactactt actctagctt cccggcaaca attaatagac tggatggagg cggataaagt 4380
     tgcaggacca cttctgcgct cggcccttcc ggctggctgg tttattgctg ataaatctgg 4440
     agcoggtgag cgtgggtctc gcggtatcat tgcagcactg gggccagatg gtaagccctc 4500
     ccgtatcgta gttatctaca cgacggggag tcaggcaact atggatgaac gaaatagaca 4560
15
     gategetgag ataggtgeet caetgattaa geattggtaa etgteagace aagtttaete 4620
     atatatactt tagattgatt taaaacttca tttttaattt aaaaggatct aggtgaagat 4680
     cctttttgat aatctcatga ccaaaatccc ttaacgtgag ttttcgttcc actgagcgtc 4740
     agaccccgta gaaaagatca aaggatcttc ttgagatcct ttttttctgc gcgtaatctg 4800
     ctgcttgcaa acaaaaaac caccgctacc agcggtggtt tgtttgccgg atcaagagct 4860
20
     accaactett ttteegaagg taactggett cagcagageg cagataceaa atactgteet 4920
     tctagtgtag ccgtagttag gccaccactt caagaactct gtagcaccgc ctacatacct 4980
     cgctctgcta atcctgttac cagtggctgc tgccagtggc gataagtcgt gtcttaccgg 5040
     gttggactca agacgatagt tacggataag gcgcagcggt cgggctgaac ggggggttcg 5100
     tgcacacage ccagettgga gegaacgace tacacegaae tgagataeet acagegtgag 5160
25
     ctatgagaaa gcgccacgct tcccgaaggg agaaaggcgg acaggtatcc ggtaagcggc 5220
     agggteggaa caggagageg caegagggag ettecagggg gaaacgeetg gtatettat 5280 agteetgteg ggtttegeea cetetgaett gagegtegat ttttgtgatg etegteaggg 5340
     gggcggagcc tatggaaaaa cgccagcaac gcggcctttt tacggttcct ggccttttgc 5400
     tggccttttg ctcacatgtt ctttcctgcg ttatcccctg attctgtgga taaccgtatt 5460
30
     accgcctttg agtgagctga taccgctcgc cgcagccgaa cgaccgagcg cagcgagtca 5520
     gtgagcgagg aagcggaaga gcgcccaata cgcaaaccgc ctctccccgc gcgttggccg 5580
     atteattaat geagetggea egacaggttt ceegactgga aagegggeag tgagegeae 5640
     gcaattaatg tgagttaget cactcattag gcaceccagg etttacaett tatgetteeg 5700
35
     gctcgtatgť tgtgtggaat tgtgagcgga ťaacaattťc acacaggaaa cagctatgac 5760
     catgattacg ccaagetete tagageteta gagetetaga getetagaga gettgeatge 5820
     cggggtacca gettegtage tagaacatea tgttetggga tateagette gtagetagaa 5880
     catcatgttc tggtaccccg gtcga
                                                                         5905
40
     <210> 14
     <211> 6052
     <212> DNA
     <213> Artificial Sequence
45.
     <220>
    <223> Description of Artificial Sequence: vector pTGFG64
     <400> 14
50
    cgcgttgaca ttgattattg actagttatt aatagtaatc aattacgggg tcattagttc 60
    atageceata tatggagtte egegttacat aacttaeggt aaatggeeeg eetggetgae 120
    cgcccaacga cccccgccca tigacgtcaa taatgacgta tgttcccata gtaacgccaa 180
    tagggactit ccattgacgt caatgggtgg agtatttacg gtaaactgcc cacttggcag 240
    tacatcaagt gtatcatatg ccaagtacgc cccctattga cgtcaatgac ggtaaatggc 300
55
    ccgcctggca ttatgcccag tacatgacct tatgggactt tectacttgg cagtacatet 360
    acgtattagt categoratt accatggtga tgeggttttg geagtacate aatgggegtg 420
    gatageggtt tgacteaegg ggattteeaa gteteeacee eattgaegte aatgggagtt 480
    tgttttggca ccaaaatcaa cgggactttc caaaatgtcg taacaactcc gccccattga 540
    cgcaaatggg cggtaggcgt gtacggtggg aggtctatat aagcagagct ctctggctaa 600
   ctagagaacc cactgettac tggettateg aaattaatac gaeteactat agggagaece 660
    aagcttgcat gccaattccg caaaggttat gcagcgcgtg aacatgatca tggcagaatc 720
    accaggeete ateaceatet geetttagg atatetaete agtgetgaat gtacagtttt 780
```

tcttgatcat gaaaacgcca acaaaattct gaatcggcca aagaggtata attcaggtaa 840 attggaagag tttgttcaag ggaaccttga gagagaatgt atggaagaaa agtgtagttt 900 tgaagaagca cgagaagtti tigaaaacac tgaaagaaca actgaatttt ggaagcagta 960 tgttgatgga gatcagtgtg agtccaatcc atgtttaaat ggcggcagtt gcaaggatga 1020 cattaattee tatgaatgtt ggtgteeett tggatttgaa ggaaagaaet gtgaattaga 1080 5 tgtaacatgt aacattaaga atggcagatg cgagcagttt tgtaaaaata gtgctgataa 1140 caaggtggtt tgctcctgta ctgagggata tcgacttgca gaaaaccaga agtcctgtga 1200 accagcagtg ccatttccat gtggaagagt ttctgtttca caaacttcta agctcacccg 1260 tgctgagact gtttttcctg atgtggacta tgtaaattct actgaagctg aaaccatttt 1320 10 ggataacatc actcaaagca cccaatcatt taatgacttc actcgggttg ttggtggaga 1380 agatgecaaa ecaggteaat teeettggea ggttgttttg aatggtaaag ttgatgeatt 1440 ctgtggagge tetategtta atgaaaatg gattgtaact getgeecact gtgttgaaac 1500 tggtgttaaa attacagttg tcgcaggtga acataatatt gaggagacag aacatacaga 1560 gcaaaagcga aatgtgatto gaattattoo toaccacac tacaatgcag ctattaataa 1620 15 gtacaaccat gacattgeee ttetggaact ggaegaacee ttagtgetaa acagetaegt 1680 tacacctatt tgcattgctg acaaggaata cacgaacatc ttcctcaaat ttggatctgg 1740 ctatgtaagt ggctggggaa gagtcttcca caaagggaga tcagctttag ttcttcagta 1800 ccttagagtt ccactigttg accgagecae atgtettega tetacaaagt teaccateta 1860 taacaacatg ttctgtgctg gcttccatga aggaggtaga gattcatgtc aaggagatag 1920 tgggggaccc catgttactg aagtggaagg gaccagtttc ttaactggaa ttattagctg 1980 20 gggtgaagag tgtgcaatga aaggcaaata tggaatatat accaaggtat cccggtatgt 2040 caactggatt aaggaaaaa caaagctcac ttaatgggat cggtcgagcg gccgcgactc 2100 tactagagga tctttgtgaa ggaaccttac ttctgtggtg tgacataatt ggacaaacta 2160 cctacagaga tttaaagctc taaggtaaat ataaaatttt taagtgtata atgtgttaaa 2220 ctactgattc taattgtttg tgtattttag attccaacct atggaactga tgaatgggag 2280 25 cagtggtgga atgcctttaa tgaggaaaac ctgttttgct cagaagaaat gccatctagt 2340 gatgatgagg ctactgctga ctctcaacat tctactcctc caaaaaagaa gagaaaggta 2400 gaagacccca aggactttcc ttcagaattg ctaagttttt tgagtcatgc tgtgtttagt 2460 aatagaacte tigettgett tgetatttae accaeaagg aaaaagetge actgetatae 2520 aagaaaatta tggaaaaata ttctgtaacc tttataagta ggcataacag ttataatcat 2580 30 aacatactgt tittettac tecacacagg catagagigt cigetattaa taactatget 2640 caaaaattgt gtacctttag ctttttaatt tgtaaagggg ttaataagga atatttgatg 2700 tatagtgcct tgactagaga tcataatcag ccataccaca tttgtagagg ttttacttgc 2760 tttaaaaaac ctcccacacc tcccctgaa cctgaaacat aaaatgaatg caattgttgt 2820 35 tgttaacttg tttattgcag cttataatgg ttacaaataa agcaatagca tcacaaattt 2880 cacaaataaa gcatttttt cactgcattc tagttgtggt ttgtccaaac tcatcaatgt 2940 atcttatcat gtctggatcc ccggggggta ccagcttcgt agctagaaca tcatgttctg 3000 ggatatcage ttegtageta gaacatcatg ttetggtace ceetetaga gegaattaat 3060 tcactggccg tcgttttaca acgtcgtgac tgggaaaacc ctggcgttac ccaacttaat 3120 cgccttgcag cacatecece tttcgccage tggcgtaata gegaagagge cegcacegat 3180 40 cgcccttccc aacagttgcg cagcctgaat ggcgaatggc ggggtaccag cttcgtagct 3240 agaacatcat gttctgggat atcagcttcg tagctagaac atcatgttct ggtaccccgc 3300 ctgatgcggt attttctcct tacgcatctg tgcggtattt cacaccgcat atggtgcact 3360 ctcagtacaa tetgetetga tgccgcatag ttaagccage ceegacaeee gccaacaeee 3420 45 gctgacgcgc cctgacgggc ttgtctgctc ccggcatccg cttacagaca agctgtgacc 3480 gtotocggga gotgoatgtg toagaggttt toacogtoat cacegaaacg cgcgagacga 3540 aagggcacca gaacatgatg ttctagctac gaagctgata tcccagaaca tgatgttcta 3600 gctacgaage tggtaceceg cetegígata egeétaíttt tataggttaa tgteatgata 3660 ataatggttt cttagacgtc aggtggcact tttcggggaa atgtgcgcgg aacccctatt 3720 50 tgtttatttt tctaaataca ttcaaatatg tatccgctca tgagacaata accctgataa 3780 atgcttcaat aatattgaaa aaggaagagt atgagtattc aacatttccg tgtcgccctt 3840 attecetttt ttgeggeatt ttgeetteet gtitttgete acceagaaac getggtgaaa 3900 gtaaaagatg ctgaagatca gttgggtgca cgagtgggtt acatcgaact ggatctcaac 3960 agcggtaaga tccttgagag ttttcgcccc gaagaacgtt ttccaatgat gagcactttt 4020 55 aaagttetge tatgtggege ggtattatee egtattgaeg eegggeaaga geaacteggt 4080 cgccgcatac actattctca gaatgacttg gttgagtact caccagtcac agaaaagcat 4140 cttacggatg gcatgacagt aagagaatta tgcagtgctg ccataaccat gagtgataac 4200 actgeggeca acttacttet gacaacgate ggaggaeega aggagetaae egetittetg 4260 cacaacatgg gggatcatgt aactcgcctt gatcgttggg aaccggagct gaatgaagcc 4320 60 ataccaaacg acgagegtga caccacgatg cetgtageaa tggcaacaac gttgegeaaa 4380 ctattaactg gcgaactact tactctagct tcccggcaac aattaataga ctggatggag 4440 geggataaag ttgeaggace acttetgege teggeeette eggetggetg gtttattget 4500

```
gataaatctg gagccggtga gcgtgggtct cgcggtatca ttgcagcact ggggccagat 4560
     ggtaagccct cccgtatcgt agttatctac acgacgggga gtcaggcaac tatggatgaa 4620
     cgaaatagac agategetga gataggtgee teaetgatta ageattggta actgteagae 4680
     caagtttact catatact ttagattgat ttaaaacttc attttaatt taaaaggatc 4740
     taggtgaaga teettttga taateteatg accaaaatee ettaaegtga gttttegtte 4800
     cactgagegt cagacceegt agaaaagate aaaggatett ettgagatee ttttttetg 4860
     cgcgťaátčt gcťgcttgča aácaaaaaa ccaccgctac cagcggtggt ttgtttgccg 4920
     gatcaagage taccaactet tttteegaag gtaactgget teageagage geagatacea 4980
     aatactgtcc ttctagtgta gccgtagtta ggccaccact tcaagaactc tgtagcaccg 5040
     cctacatacc tegetetget aatectgtta ccagtggetg etgecagtgg egataagteg 5100
10
     tgtcttaccg ggttggactc aagacgatag ttaccggata aggcgcagcg gtcgggctga 5160
     acggggggtt cgtgcacaca gcccagcttg gagcgaacga cctacaccga actgagatac 5220
     ctacagegtg agetatgaga aagegeeaeg etteeegaag ggagaaagge ggacaggtat 5280
     ceggtaageg geagggtegg aacaggagag egeacgaggg agetteeagg gggaaaegee 5340
     tggtatettt atagteetgt egggtítege caectetgae tígagegteg atitttgiga 5400
15
     tgctcgtcag ggggggggag cctatggaaa aacgccagca acgcggcctt tttacggttc 5460
     ctggcctttt gctggccttt tgctcacatg ttctttcctg cgttatcccc tgattctgtg 5520
     gataaccgta ttaccgcctt tgagtgagct gataccgctc gccgcagccg aacgaccgag 5580
     cgcagcgagt cagtgagcga ggggtaccag aacatgatgt tctagctacg aagctgatat 5640
20
     cccagaacat gatgttctag ctacgaagct ggtaccccag cggaagagcg cccaatacgc 5700
     aaaccgcctc tccccgcgcg ttggccgatt cattaatgca gctggcacga caggtttccc 5760
     gactggaaag cgggcagtga gcgcaacgca attaatgtga gttagctcac tcattaggca 5820
     ccccaggett tacactttat getteegget egtatgttgt gtggaattgt gageggataa 5880
     caatttcaca caggaaacag ctatgaccat gattacgcca agctctctag agctctagag 5940
25
     ctctagagct ctagagagct tgcatgccgg ggtaccagct tcgtagctag aacatcatgt 6000
     tctgggatat cagcttcgta gctagaacat catgttctgg taccccggtc ga
     <210> 15
30
     <211> 4344
     <212> DNA
     <213> Artificial Sequence
     <220>
     <223> Description of Artificial Sequence: vector pTGFG67
35
     <400> 15
     cgcgttgaca ttgattattg actagttatt aatagtaatc aattacgggg tcattagttc 60
     atageceata tatggagtte egegttaeat aacttaeggt aaatggeeeg eetggetgae 120
40
     cgcccaacga cccccgccca ttgacgtcaa taatgacgta tgttcccata gtaacgccaa 180
     tagggactit ccattgacgt caatgggtgg agtatttacg gtaaactgcc cacttggcag 240
     tacatcaagt gtatcatatg ccaagtacgc cccctattga cgtcaatgac ggtaaatggc 300
     cegeetggea ttatgeceag tacatgacet tatgggaett teetaettgg cagtacatet 360
     acgtattagt categetatt accatggtga tgeggttttg geagtacate aatgggegtg 420
    gatageggtt tgactcaegg ggatttecaa gtetecacee cattgaegte aatgggagtt 480
45
     tgttttggca ccaaaatcaa cgggactttc caaaatgtcg taacaactcc gccccattga 540
    cgcaaatggg cggtaggcgt gtacggtggg aggtctatat aagcagagct ctctggctaa 600
    ctagagaacc cactgcttac tggcttatcg aaattaatac gactcactat agggagaccc 660
    aagettgace tegageaage ggeegegact etactagagg atetttgtga aggaacetta 720
50
    cttctgtggt gtgacataat tggacaaact acctacagag atttaaagct ctaaggtaaa 780
    tataaaattt ttaagtgtat aatgtgttaa actactgatt ctaattgttt gtgtatttta 840
    gattccaacc tatggaactg atgaatggga gcagtggtgg aatgccttta atgaggaaaa 900
    cctgttttgc tcagaagaaa tgccatctag tgatgatgag gctactgctg actctcaaca 960 ttctactcct ccaaaaaaga agagaaaggt agaagacccc aaggactttc cttcagaatt 1020
55
    gctaagtttt ttgagtcatg ctgtgtttag taatagaact cttgcttgct ttgctattta 1080
    caccacaaag gaaaaagctg cactgctata caagaaaatt atggaaaaat attctgtaac 1140
    ctttataagt aggcataaca gttataatca taacatactg ttttttctta ctccacacag 1200
    gcatagagtg tetgetatta ataactatge teaaaaattg tgtaeettta getttttaat 1260
    ttgtaaaggg gttaataagg aatatttgat gtatagtgcc ttgactagag atcataatca 1320
60
    gecataccae atttgtagag gttttacttg etttaaaaaa eeteecaeae eteecetga 1380
    acctgaaaca taaaatgaat gcaattgttg ttgttaactt gtttattgca gcttataatg 1440
    gttacaaata aagcaatagc atcacaaatt tcacaaataa agcatttitt tcactgcatt 1500
```

(A)

```
ctagttgtgg tttgtccaaa ctcatcaatg tatcttatca tgtctggatc cccgggtacc 1560
        ctctagagcg aattaattca ctggccgtcg ttttacaacg tcgtgactgg gaaaaccctg 1620
         gcgttaccca acttaatcgc cttgcagcac atccccttt cgccagctgg cgtaatagcg 1680
        aagaggeeeg cacegatege eetteecaae agttgegeag eetgaatgge gaatggegee 1740
        tgatgcggta ttttctcctt acgcatctgt gcggtatttc acaccgcata tggtgcactc 1800
        tragtaria tragetetgat gregoatagt taagerage regaracer coaacacer 1860 etgacgege regarget tyletgeter regeatere thanks the second of the second regarder regarde
        tctccgggag ctgcatgtgt cagaggtttt caccgtcatc accgaaacgc gcgagacgaa 1980
        agggggggta ccagettegt agetagaaca teatgttetg ggatateage ttegtageta 2040
        gaacatcatg ttctggtacc cccctcgtga tacgcctatt tttataggtt aatgtcatga 2100
 10
        taataatggt ttettagaeg teaggtggea ettttegggg aaatgtgege ggaaceeta 2160
        tttgtttatt tttctaaata cattcaaata tgtatccgct catgagacaa taaccctgat 2220
        aaatgettea ataatattga aaaaggaaga gtatgagtat teaacattte egtgtegeee 2280
        ttattccctt ttttgcggca ttttgccttc ctgttttgc tcacccagaa acgctggtga 2340
        aagtaaaaga tgctgaagat cagttgggtg cacgagtggg ttacatcgaa ctggatctca 2400
 15
        acageggtaa gateettgag agittiegee eegaagaaeg ttttecaatg atgageaett 2460
        ttaaagttct gctatgtggc gcggtattat cccgtattga cgccgggcaa gagcaactcg 2520
        gtogoogcat acactattot cagaatgact tggttgagta ctcaccagtc acagaaaago 2580
        atcttacgga tggcatgaca gtaagagaat tatgcagtgc tgccataacc atgagtgata 2640
        acactgogge caacttaett otgacaacga toggaggaco gaaggageta accgottttt 2700
 20
        tgcacaacat gggggatcat gtaactcgcc ttgatcgttg ggaaccggag ctgaatgaag 2760
        ccataccaaa cgacgagcgt gacaccacga tgcctgtagc aatggcaaca acgttgcgca 2820
        aactattaac tggcgaacta cttactctag cttcccggca acaattaata gactggatgg 2880
        aggoggataa agttgcagga ccacttotgo gotoggooot tooggotggo tggtttattg 2940
        ctgataaatc tggagccggt gagcgtgggt ctcgcggtat cattgcagca ctggggccag 3000
25
        atggtaagce ctcccgtate gtagttatet acacgacggg gagtcaggca actatggatg 3060
        aacgaaatag acagateget gagataggtg ceteaetgat taagcattgg taactgteag 3120
        accaagttta ctcatatata ctttagattg atttaaaact tcatttttaa tttaaaagga 3180
        tctaggtgaa gatcctttt gataatctca tgaccaaaat cccttaacgt gagttttcgt 3240
        tccactgage gtcagacece gtagaaaaga tcaaaggate ttettgagat cetttttte 3300
30
        tgcgcgtaat ctgctgcttg caaacaaaaa aaccaccgct accagcggtg gtttgtttgc 3360
        cggatcaaga gctaccaact ctttttccga aggtaactgg cttcagcaga gcgcagatac 3420
        caaatactgt tettetagtg tageegtagt taggecacca etteaagaae tetgtageae 3480
       cgcctacata cctcgctctg ctaatcctgt taccagtggc tgctgccagt ggcgataagt 3540
35
       cgtgtettac egggttggac teaagacgat agttacegga taaggegeag eggteggget
       gaacgggggg třegtgeaca cagcecaget tggagegaac gacetacace gaactgagat
       acctacagcg tgagctatga gaaagcgcca cgcttcccga agggagaaag gcggacaggt
       atccggtaag cggcagggtc ggaacaggag agcgcacgag ggagcttcca gggggaaacg 3780
       cctggtatct ttatagtect gtegggttte gecacctetg acttgagegt egatttttgt 3840
40
       gatgetegte aggggggegg ageetatgga aaaacgecag caacgeggee titttaeggt 3900
       teetggeett ttgetggeet tttgeteaca tgttetttee tgegttatee eetgattetg 3960
       tggataaccg tattaccgcc tttgagtgag ctgataccgc tcgccgcagc cgaacgaccg 4020
       agegeagega gteagtgage gaggaagegg aagagegeee aataegeaaa eegeetetee 4080
       ccgcgcgttg gccgattcat taatgcagct ggcacgacag gtttcccgac tggaaagcgg 4140
45
       gcagtgagcg caacgcaatt aatgtgagtt agctcactca ttaggcaccc caggctttac 4200
       actttatget teeggetegt atgttgtgtg gaattgtgag eggataacaa tttcacacag 4260
       gaaacageta tgaccatgat taegecaage tetetagage tetagagete tagageteta 4320
       gagagettge atgeetgeag gteg
50
       <210> 16
       <211> 4496
       <212> DNA
       <213> Artificial Sequence
55
       <223> Description of Artificial Sequence: vector pTGFG82
       <400> 16
60
       cgcgttgaca ttgattattg actagttatt aatagtaatc aattacgggg tcattagttc 60
       atageceata tatggagtte egegttacat aacttaeggt aaatggeeeg eetggetgae 120
       cgcccaacga cccccgccca ttgacgtcaa taatgacgta tgttcccata gtaacgccaa 180
```

tagggacttt ccattgacgt caatgggtgg agtatttacg gtaaactgcc cacttggcag 240 tacatcaagt gtatcatatg ccaagtacgc cccctattga cgtcaatgac ggtaaatggc 300 ecgeetggea tratgeceag tacatgacet tatgggactt teetaettgg cagtacatet 360 acgtattagt categetatt accatggtga tgeggttttg geagtacate aatgggegtg 420 gatagcggtt tgactcacgg ggatttccaa gtctccaccc cattgacgtc aatgggagtt 480 5 tgttttggca ccaaaatcaa cgggactttc caaaatgtcg taacaactcc gcccattga 540 cgcaaatggg cggtaggcgt gtacggtggg aggtctatat aagcagagct ctctggctaa 600 ctagagaacc cactgcttac tggcttatcg aaattaatac gactcactat agggagaccc 660 aagettgace tegageaage ggeegegact etactagagg atetttgtga aggaacetta 720 cttctgtggt gtgacataat tggacaaact acctacagag atttaaagct ctaaggtaaa 780 10 tataaaattt ttaagtgtat aatgtgttaa actactgatt ctaattgttt gtgtatttta 840 gattccaacc tatggaactg atgaatggga gcagtggtgg aatgccttta atgaggaaaa 900 cctgttttgc tcagaagaaa tgccatctag tgatgatgag gctactgctg actctcaaca 960 ttetacteet ecaaaaaaga agagaaaggt agaagaeeee aaggaettte etteagaatt 1020 gctaagtttt ttgagtcatg ctgtgtttag taatagaact cttgcttgct ttgctattta 1080 15 caccacaaag gaaaaagctg cactgctata caagaaaatt atggaaaaat attctgtaac 1140 ctttataagt aggcataaca gttataatca taacatactg ttttttctta ctccacacag 1200 gcatagagtg tetgetatta ataactatge teaaaaattg tgtacettta getttttaat 1260 ttgtaaaggg gttaataagg aatatttgat gtatagtgcc ttgactagag atcataatca 1320 gecataceae atttgtagag gttttaettg etttaaaaaa eeteecaeae eteeceetga 1380 20 acctgaaaca taaaatgaat gcaattgttg ttgttaactt gtttattgca gcttataatg 1440 gttacaaata aagcaatagc atcacaaatt tcacaaataa agcattttt tcactgcatt 1500 ctagttgtgg tttgtccaaa ctcatcaatg tatcttatca tgtctggatc cccggggggt 1560 accagetteg tagetagaac atcatgttet gggatateag ettegtaget agaacateat 1620 25 gttctggtac cccctctag agcgaattaa ttcactggcc gtcgttttac aacgtcgtga 1680 ctgggaaaac cctggcgtta cccaacttaa tcgccttgca gcacatcccc ctttcgccag 1740 ctggcgtaat agcgaagagg cccgcaccga tcgcccttcc caacagttgc gcagcctgaa 1800 tggcgaatgg cgcctgatgc ggtattttct ccttacgcat ctgtgcggta tttcacaccg 1860 catatggtgc actctcagta caatctgctc tgatgccgca tagttaagcc agccccgaca 1920 30 cccgccaaca cccgctgacg cgccctgacg ggcttgtctg ctcccggcat ccgcttacag 1980 acaagetgtg accettece ggagetgeat gtgteagagg tttteacegt cateacegaa 2040 acgcgcgaga cgaaagggcg gggtaccaga acatgatgtt ctagctacga agctgatatc 2100 ccagaacatg atgitctage tacgaagetg gtacceegge ctegtgatae geetattitt 2160 ataggttaat gtcatgataa taatggtttc ttagacgtca ggtggcactt ttcggggaaa 2220 tgtgcgcgga accectattt gtttatttt ctaaatacat tcaaatatgt atccgctcat 2280 35 gagacaataa ccctgataaa tgcttcaata atattgaaaa aggaagagta tgagtattca 2340 acattteegt gtegeeetta tiecettttt tgeggeattt tgeetieetg tititgetea 2400 cccagaaacg ctggtgaaag taaaagatgc tgaagatcag ttgggtgcac gagtgggtta 2460 categaactg gateteaaca geggtaagat cettgagagt tttegeeeeg aagaacgttt 2520 40 tccaatgatg agcactttta aagttctgct atgtggcgcg gtattatccc gtattgacgc 2580 cgggcaagag caactcggtc gccgcataca ctattctcag aatgacttgg ttgagtactc 2640 accagtcaca gaaaagcate ttacggatgg catgacagta agagaattat gcagtgctgc 2700 cataaccatg agtgataaca ctgcggccaa cttacttctg acaacgatcg gaggaccgaa 2760 ggagctaacc gcttttttgc acaacatggg ggatcatgta actcgccttg atcgttggga 2820 accggagetg aatgaageca taccaaacga cgagegtgae accaegatge etgtageaat 2880 45 ggcaacaacg ttgcgcaaac tattaactgg cgaactactt actctagctt eccggcaaca 2940 attaatagac tggatggagg cggataaagt tgcaggacca cttctgcgct cggcccttcc 3000 ggctggctgg tttattgctg ataaatctgg agccggtgag cgtgggtctc gcggtatcat 3060 tgcagcactg gggccagatg gtaagccctc ccgtatcgta gttatctaca cgacggggag 3120 tcaggcaact atggatgaac gaaatagaca gatcgctgag ataggtgcct cactgattaa 3180 50 gcattggtaa ctgtcagacc aagtttactc atatatactt tagattgatt taaaacttca 3240 tttttaattt aaaaggatct aggtgaagat cetttttgat aateteatga eeaaaateee 3300 ttaacgtgag ttttcgttcc actgagcgtc agaccccgta gaaaagatca aaggatcttc 3360 ttgagateet tttttetge gegtaatetg etgettgeaa acaaaaaac cacegetace 3420 55 ageggtggtt tgtttgeegg ateaagaget aceaactett ttteegaagg taactggett 3480 cagcagageg cagataccaa atactgteet tetagtgtag cegtagttag gecaccaett 3540 caagaactet gtagcaccge ctacatacet egetétgeta atéctgttac cagtggetge 3600 tgccagtggc gataagtcgt gtcttaccgg gttggactca agacgatagt taccggataa 3660 ggcgcagcgg tcgggctgaa cggggggttc gtgcacacag cccagcttgg agcgaacgac 3720 60 ctacaccgaa ctgagatacc tacagcgtga gctatgagaa agcgccacgc ttcccgaagg 3780 gagaaaggcg gacaggtatc cggtaagcgg cagggtcgga acaggagagc gcacgaggga 3840 gettecaggg ggaaacgeet ggtatettta tagteetgte gggtttegee acetetgaet 3900

```
tgagcgtcga tttttgtgat gctcgtcagg ggggcggagc ctatggaaaa acgccagcaa 3960
     egeggeettt ttaeggttee tggeetttig etggeettti geteacatgt tettteetge 4020
     gttatcccct gattetgtgg ataaccgtat taccgccttt gagtgagetg ataccgctcg 4080
     ccgcagccga acgaccgagc gcagcgagtc agtgagcgag gaagcggaag agcgcccaat 4140
     acgcaaaccg ceteteceg egegttggee gatteattaa tgcagetgge acgacaggtt 4200 teecgaetgg aaagegggea gtgagegeaa egeaattaat gtgagttage teacteatta 4260
     ggcacccag gctttacact ttatgcttcc ggctcgtatg ttgtgggaa ttgtgagcgg 4320
     ataacaattt cacacaggaa acagctatga ccatgattac gccaagctct ctagagctct 4380
     agagetetag agetetagag agettgeatg eeggggtace agettegtag etagaacate 4440
10
     atgitciggg atatcagcit cgtagctaga acatcatgit ciggtacccc ggtcga
     <210> 17
     <211> 4644
15
     <212> DNA
     <213> Artificial Sequence
     <220>
     <223> Description of Artificial Sequence: vector pTGFG95
20
     <400> 17
     cgcgttgaca ttgattattg actagttatt aatagtaatc aattacgggg tcattagttc 60
     atageceata tatggagtte egegttaeat aacttaeggt aaatggeeeg eetggetgae 120
     egeccaaega eccegecca tigaegteaa taatgaegta tgtteecata gtaaegecaa 180
     tagggactit ccattgacgt caatgggtgg agtatttacg gtaaactgcc cacttggcag 240
25
     tacatcaagt gtatcatatg ccaagtacgc cccctattga cgtcaatgac ggtaaatggc 300
     ccgcctggca ttatgcccag tacatgacct tatgggactt tcctacttgg cagtacatct 360
     acgtattagt categotatt accatggtga tgcggttttg gcagtacatc aatgggcgtg 420
     gatageggtt tgactcaegg ggatttecaa gtetecaece cattgaegte aatgggagtt 480
     tgttttggca ccaaaatcaa cgggactttc caaaatgtcg taacaactcc gccccattga 540
30
     cgcaaatggg cggtaggcgt gtacggtggg aggtctatat aagcagagct ctctggctaa 600
     ctagagaacc cactgettae tggettateg aaattaatae gaeteaetat agggagaeee 660
     aagettgace tegageaage ggeegegact etactagagg atetttgtga aggaacetta 720
     cttctgtggt gtgacataat tggacaaact acctacagag atttaaagct ctaaggtaaa 780
35
     tataaaattt ttaagtgtat aatgtgttaa actactgatt ctaattgttt gtgtatttta 840
     gattccaacc tatggaactg atgaatggga gcagtggtgg aatgccttta atgaggaaaa 900
     cctgttttgc tcagaagaaa tgccatctag tgatgatgag gctactgctg actctcaaca 960
     ttctactcct ccaaaaaaga agagaaaggť agaagacccc aaggactttc cttcagaatt 1020
     gctaagtttt ttgagtcatg ctgtgtttag taatagaact cttgcttgct ttgctattta 1080
     caccacaag gaaaaagctg cactgctata caagaaaatt atggaaaaat attctgtaac 1140
40
    ctttataagt aggcataaca gttataatca taacatactg ttttttctta ctccacacag 1200
     gcatagagig toigctatta ataactatgo toaaaaattg tgtaccttta gctttttaat 1260
     ttgtaaaggg gttaataagg aatatttgat gtatagtgcc ttgactagag atcataatca 1320
    gecataccac atttgtagag gttttacttg ctttaaaaaa cctcccacac ctcccctga 1380
45
     acctgaaaca taaaatgaat gcaattgttg ttgttaactt gtttattgca gcttataatg 1440
    gttacaaata aagcaatagc atcacaaatt tcacaaataa agcatttitt tcactgcati 1500
    ctagttgtgg tttgtccaaa ctcatcaatg tatcttatca tgtctggatc cccggggggt 1560
    accagetteg tagetagaac atcatgttet gggatateag ettegtaget agaacateat 1620
    gttctggtac ccccctctag agcgaattaa ttcactggcc gtcgttttac aacgtcgtga 1680
50
    ctgggaaaac cctggcgtta cccaacttaa tcgccttgca gcacatcccc ctttcgccag 1740
    ctggcgtaat agcgaagagg cccgcaccga tcgcccttcc caacagttgc gcagcctgaa 1800
    tggcgaatgg cggggtacca gcttcgtagc tagaacatca tgttctggga tatcagcttc 1860
    gtagctagaa catcatgttc tggtaccccg cctgatgcgg tattttctcc ttacgcatct 1920
    gtgcggtatt tcacaccgca tatggtgcac tctcagtaca atctgctctg atgccgcata 1980
    gttaagccag coccgacace cgccaacace cgctgacgcg ccctgacggg cttgtctgct 2040
55
    cccggcatcc gcttacagac aagctgtgac cgtctccggg agctgcatgt gtcagaggtt 2100
    ttcaccgtca tcaccgaaac gcgcgagacg aaagggctac cagaacatga tgttctagct 2160
    acgaagctga tatcccagaa catgatgttc tagctacgaa gctggtaccc cgcctcgtga 2220
    tacgcctatt tttataggtt aatgtcatga taataatggt ttcttagacg tcaggtggca 2280
60
    cttttcgggg aaatgtgcgc ggaaccccta tttgtttatt tttctaaata cattcaaata 2340
    tgtatccgct catgagacaa taaccctgat aaatgcttca ataatattga aaaaggaaga 2400
    gtatgagtat tcaacatttc cgtgtcgccc ttattccctt ttttgcggca ttttgccttc 2460
```

```
ctgtttttgc tcacccagaa acgctggtga aagtaaaaga tgctgaagat cagttgggtg 2520
      cacgagtggg ttacatcgaa ctggatctca acageggtaa gateettgag agttttegee 2580
      ccgaagaacg ttttccaatg atgagcactt ttaaagttct gctatgtggc gcggtattat 2640
      cccgtattga cgccgggcaa gagcaactcg gtcgccgcat acactattct cagaatgact 2700
      tggttgagta ctcaccagtc acagaaaagc atcttacgga tggcatgaca gtaagagaat 2760
      tatgcagtgc tgccataacc atgagtgata acactgcggc caacttactt ctgacaacga 2820
      tcggaggacc gaaggagcta accgcttttt tgcacaacat gggggatcat gtaactcgcc 2880
      ttgatcgttg ggaaccggag ctgaatgaag ccataccaaa cgacgagcgt gacaccacga 2940
      tgcctgtagc aatggcaaca acgttgcgca aactattaac tggcgaacta cttactctag 3000
 10
      cttcccggca acaattaata gactggatgg aggcggataa agttgcagga ccacttctgc 3060
     geteggeet teeggetgge tggtttattg etgataaate tggageeggt gagegtgggt 3120
      ctcgcggtat cattgcagca ctggggccag atggtaagcc ctcccgtatc gtagttatct 3180
      acacgacggg gagtcaggca actatggatg aacgaaatag acagatcgct gagataggtg 3240
      cctcactgat taagcattgg taactgtcag accaagttta ctcatatata ctttagattg 3300
 15
      atttaaaact teatttttaa tttaaaagga tetaggtgaa gateettttt gataatetea 3360
      tgaccaaaat cccttaacgt gagttttcgt tccactgagc gtcagacccc gtagaaaaga 3420
      tcaaaggatc ttcttgagat ccttttttc tgcgcgtaat ctgctgcttg caaacaaaaa 3480
      aaccaccgct accagcggtg gtttgtttgc cggatcaaga gctaccaact ctttttccga 3540
      aggtaactgg cttcagcaga gcgcagatac caaatactgt ccttctagtg tagccgtagt 3600
 20
      taggecacca etteaagaae tetgtageae egeetacata eetegetetg etaateetgt 3660
      taccagtggc tgctgccagt ggcgataagt cgtgtcttac cgggttggac tcaagacgat 3720
      agttacegga taaggegeag eggteggget gaaegggggg ttegtgeaca eageceaget 3780
      tggagcgaac gacctacacc gaactgagat acctacagcg tgagctatga gaaagcgcca 3840
      cgcttcccga agggagaaag gcggacaggt atccggtaag cggcagggtc ggaacaggag 3900
 25
      agcgcacgag ggagcttcca gggggaaacg cctggtatct ttatagtcct gtcgggtttc 3960
      gecaectetg aettgagegt egatttttgt gatgetegte aggggggegg agectatgga 4020
      aaaacgccag caacgcggcc tttttacggt tcctggcctt ttgctggcct tttgctcaca 4080
      tgttetttee tgegttatee eetgattetg tggataaceg tattacegee tttgagtgag 4140
      ctgataccge tegeegeage egaacgaceg agegeagega gteagtgage gaggggtace 4200 agaacatgat gttetageta egaagetgat ateccagaae atgatgttet agetacgaag 4260
 30
      ctggtacccc agcggaagag cgcccaatac gcaaaccgcc tctccccgcg cgttggccga 4320
      ttcattaatg cagctggcac gacaggtttc ccgactggaa agcgggcagt gagcgcaacg 4380
      caattaatgt gagttagete acteattagg caceecagge tttacaettt atgetteegg 4440
      ctcgtatgtt gtgtggaatt gtgagcggat aacaatttca cacaggaaac agctatgacc 4500
 35
      atgattacgc caagetetet agagetetag agetetagag etetagagag ettgeatgee 4560
      ggggtaccag cttcgtagct agaacatcat gttctgggat atcagcttcg tagctagaac 4620
      atcatgttct ggtaccccgg tcga
 40
      <210> 18
      <211> 933
      <212> PRT
      <213> Homo sapiens
      <400> 18
45
      Met Thr Glu Leu Lys Ala Lys Gly Pro Arg Ala Pro His Val Ala Gly
      Gly Pro Pro Ser Pro Glu Val Gly Ser Pro Leu Leu Cys Arg Pro Ala
 50
      Ala Gly Pro Phe Pro Gly Ser Gln Thr Ser Asp Thr Leu Pro Glu Val
               35
                                    40
 55
      Ser Ala Ile Pro Ile Ser Leu Asp Gly Leu Leu Phe Pro Arg Pro Cys
      Gln Gly Gln Asp Pro Ser Asp Glu Lys Thr Gln Asp Gln Gln Ser Leu
 60
      Ser Asp Val Glu Gly Ala Tyr Ser Arg Ala Glu Ala Thr Arg Gly Ala
```

	Gly	GIA	Ser	Ser 100	Ser	Ser	Pro	Pro	Glu 105	Lys	Asp	Ser	Gly	Leu 110	Leu	Asp
5	Ser	Val	Leu 115		Thr	Leu	Leu	Ala 120	Pŗo	Ser	Gly	Pro	Gly 125	Gln	Ser	Gln
10	Pro	Ser 130	Pro	Pro	Ala	Cys	Glu 135	Val ⁻	Thr	Ser	Ser	Trp 140	Cys	Leu	Phe	Gly
10	Pro 145	Glu	Leu	Pro	Glu	Asp 150	Pro	Pro	Ala	Ala	Pro 155	Ala	Thr	Gln	Arg	Val 160
15	Leu _.	Ser	Pro	Leu	Met 165	Ser	Arg	Ser	Gly	Cys 170	Lys	Val	Gly	Asp	Ser 175	Ser
	Gly	Thr	Ala	Ala 180	Ala	His	Lys	Val	Leu 185	Pro	Arg	Gly	Leu	Ser 190	Pro	Ala
20	Arg	Gln	Leu 195	Leu	Leu	Pro	Ala	Ser 200	Glu	Ser	Pro	His	Trp 205	Ser	Gly	Ala
25	Pro	Val 210	Lys	Pro	Ser	Pro	Gln 215	Ala	Ala	Ala	Val	Glu 220	Val	Glu	Glu	Glu
	Asp 225	Gly	Ser	Glu	Ser	Glu 230	Glu	Ser	Ala	Gly	Pro 235		Leu	Lys	Gly	Lys 240
30	Pro	Arg	Ala	Leu	Gly 245	Gly	Ala	Ala	Ala	Gly 250	Gly	Gly	Ala	Ala	Ala 255	Val
	Pro	Pro	Gly	Ala 260	Ala	Ala	Gly	Gly	Val 265	Ala	Leu	Val	Pro	Lys 270	Glu	Asp
35	Ser	Arg	Phe 275	Ser	Ala	Pro	Arg	Val 280	Ala	Leu	Val	Glu	Gln 285	Asp	Ala	Pro
40	Met	Ala 290	Pro	Gly	Arg	Ser	Pro 295	Leu	Ala	Thr	Thr	Val 300	Met	Asp	Phe	Il∈
40	His 305	Val	Pro	Ile	Leu	Pro 310	Leu	Asn	His	Ala	Leu 315	Leu	Ala	Ala	Arg	Thr 320
45	Arg	Gln	Leu	Leu	Glu 325	Asp	Glu	Ser		Asp 330	Gly	Gly	Ala	Gly	Ala 335	Ala
	Ser	Ala	Phe	Ala 340		Pro	Arg	Ser	Ser 345	Pro	Cys	Ala	Ser	Ser 350	Thr	Pro
50	Val	Ala	Val 355		Asp	Phe	Pro	Asp 360	Cys	Ala	Tyr	Pro	Pro 365		Ala	Glv
55	Pro	Lys 370		Asp	Ala	Tyr	Pro 375		Tyr	Ser	Asp	Phe 380		Pro	Pro	Ala
<i></i>	Leu 385	-	Ile	Lys	Glu	Glu 390	Glu	Glu	Gly	Ala	Glu 395	Ala	Ser	Ala	Arg	Se:
60	Pro	Arg	Ser	Tyr	Leu 405		Ala	Gly	Ala	Asn 410		Ala	Ala	Phe	Pro 415	

	· Pne	Pro) Leu	420	Pro	Pro	Pro	Pro	Leu 425	Pro	Pro	Arg	Ala	Thr 430		Se:
5	Arg	Pro	Gly 435	Glu	Ala	Ala	Val	Thr 440	Ala	Ala	Pro	Ala	Ser 445		Ser	Va.
10	Ser	Ser 450	Ala	Ser	Ser	Ser	Gly 455	Ser	Thr	Leu	Glu	Cys 460	Ile	Leu	Tyr	Lys
	Ala 465	Glu	Gly	Ala	Pro	Pro 470	Gln	Gln	Gly	Pro	Phe 475	Ala	Pro	Pro	Pro	Cys 480
15			Pro		403					490					495	
	Ser	Thr	Ser	Ala 500	Ser	Ala	Ala	Ala	Ala 505	Gly	Ala	Ala	Pro	Ala 510		Туг
20	Pro	Ala	Leu 515	Gly	Leu	Asn	Gly	Leu 520	Pro	Gln	Leu	Gly	Tyr 525	Gln	Ala	Ala
25	Val	Leu 530	Lys	Glu	Gly	Leu	Pro 535	Gln	Val	Tyr	Pro	Pro 540	Tyr	Leu	Asn	Туг
	Leu 545	Arg	Pro	Asp	Ser	Glu 550	Ala	Ser	Gln	Ser	Pro 555	Gln	Tyr	Ser	Phe	Glu 560
30	Ser	Leu	Pro	Gln	Lys 565	Ile	Cys	Leu	Ile	Cys 570	Gly	Asp	Glu	Ala	Ser 575	Gly
	Cys	His	Tyr	Gly 580	Val	Leu	Thr	Cys	Gly 585	Ser	Cys	Lys	Val	Phe 590	Phe	Lys
35	Arg	Ala	Met 595	Glu	Gly	Gln	His	Asn 600	Tyr	Leu	Cys	Ala	Gly 605	Arg	Asn	Asp
40	Cys	Ile 610	Val	Asp	Lys	Ile	Arg 615	Arg	Lys	Asn	Cys	Pro 620	Ala	Cys	Arg	Leu
	Arg 625	Lys	Cys	Cys	Gln	Ala 630	Gly	Met	Val	Leu	Gly 635	Gly	Arg	Lys	Phe	Lys 640
45	Lys	Phe	Asn	Lys	Val 645	Arg	Val	Val	Arg	Ala 650	Leu	Asp	Ala		Ala 655	
	Pro	Gln	Pro	Leu 660	Gly	Val	Pro	Asn	Glu 665	Ser	Gln	Ala	Leu	Ser 670	Gln	Arg
50	Phe	Thr	Phe 675	Ser	Pro	Gly	Gln	Asp 680	Ile	Gln	Leu	Ile	Pro 685	Pro	Leu	Ile
55	Asn	Leu 690	Leu	Met	Ser	Ile	Glu 695	Pro	Asp	Val	Ile	Tyr 700	Ala	Gly	His	Asp
	Asn 705	Thr	Lys	Pro	Asp	Thr 710	Ser	Ser	Ser	Leu	Leu 715	Thr	Ser	Leu	Asn	Gln 720
60	Leu	Gly	Glu	Arg	Gln 725	Leu	Leu	Ser	Val	Val 730	Lys	Trp	Ser	Lys	Ser 735	Leu

```
Pro Gly Phe Arg Asn Leu His Ile Asp Asp Gln Ile Thr Leu Ile Gln
    Tyr Ser Trp Met Ser Leu Met Val Phe Gly Leu Gly Trp Arg Ser Tyr
                                 760
    Lys His Val Ser Gly Gln Met Leu Tyr Phe Ala Pro Asp Leu Ile Leu
10
    Asn Glu Gln Arg Met Lys Glu Ser Ser Phe Tyr Ser Leu Cys Leu Thr
    785
    Met Trp Gln Ile Pro Gln Glu Phe Val Lys Leu Gln Val Ser Gln Glu
                                         810
15
    Glu Phe Leu Cys Met Lys Val Leu Leu Leu Leu Asn Thr Ile Pro Leu
                820
    Glu Gly Leu Arg Ser Gln Thr Gln Phe Glu Glu Met Arg Ser Ser Tyr
20
    Ile Arg Glu Leu Ile Lys Ala Ile Gly Leu Arg Gln Lys Gly Val Val
25
    Ser Ser Ser Gln Arg Phe Tyr Gln Leu Thr Lys Leu Leu Asp Asn Leu
                        870
                                            875
    His Asp Leu Val Lys Gln Leu His Leu Tyr Cys Leu Asn Thr Phe Ile
                    885
                                        890
                                                             895
30
    Gln Ser Arg Ala Leu Ser Val Glu Phe Pro Glu Met Met Ser Glu Val
    Ile Ala Ala Gln Leu Pro Lys Ile Leu Ala Gly Met Val Lys Pro Leu
35
    Leu Phe His Lys Lys
        930
40
    <210> 19
    <211> 2970
     <212> DNA
    <213> Homo sapiens
45
    <400> 19
    ctgaccageg cegecetece eegeceeega eecaggaggt ggagateeet eeggteeage 60
    cacattcaac acccactttc tecteetet geeectatat teeegaaace eceteetet 120
    tecettttee etecteeetg gagaeggggg aggagaaaag gggagteeag tegteatgae 180
50
    tgagetgaag geaaagggte eeegggetee ceaegtggeg ggeggeeege eeteeeega 240
    ggtcggatcc ccactgctgt gtcgcccagc cgcaggtccg ttcccgggga gccagacctc 300
    ggacacettg cetgaagttt eggecatace tateteeetg gaegggetae tetteeeteg 360
     gccctgccag ggacaggacc cctccgacga aaagacgcag gaccagcagt cgctgtcgga 420
    cgtggagggc gcatattcca gagctgaagc tacaaggggt gctggaggca gcagttctag 480
55
    teccecagaa aaggacageg gaetgetgga cagtgtettg gaeaetetgt tggegeeete 540
    aggtcccggg cagagccaac ccagccctcc cgcctgcgag gtcaccagct cttggtgcct 600
    gtttggcccc gaacttcccg aagatccacc ggctgccccc gccacccagc gggtgttgtc 660
    eccgeteatg ageeggteeg ggtgeaaggt tggagacage teegggaegg eagetgeea 720
    taaagtgctg ccccggggcc tgtcaccagc ccggcagctg ctgctcccgg cctctgagag 780
60
     ccctcactgg tccggggccc cagtgaagcc gtctccgcag gccgctgcgg tggaggttga 840
     ggaggaggat ggctctgagt ccgaggagtc tgcgggtccg cttctgaagg gcaaacctcg 900
     ggctctgggt ggcgcggcgg ctggaggagg agccgcggct gtcccgccgg gggcggcaqc 960
```

	aggaggcgtc	gccctggtcc	ссааддаада	ttcccacttc	ta2aaa		
	ggtggagcag	gacgcgccga	ccaaggaaga tggcgcccgg	acacteceea	ctagegeeca	gggtcgccct	1020
	tttcatccac	gtgcctatcc	tocctctcaa	tracgreets	ttagacacca	cggtgatgga	1080
	gctgctggaa	gacgaaagtt	acdacddcad	gaccagagact	riggeagece	gcactcggca	1140
5	gcggagttca	ccctatacct	cgtccacccc	ggtcggggtt	gedagegeet	ttgccccgcc	1200
	gtacccoccc	gacgccgagc	CCSSGGGGGGG	Cacatacact	ggcgacttcc	ccgactgcgc	1260
	gcccgctcta	aagataaagg	ccaaggacga aggaggagga	agggggggg	ccctatagcg	acttccagcc	1320
	ttcctacctt	ataaccaata	ccaaccccgc	aggegeggag	geeteegege	gctccccgcg	1380
	gcccccqctq	ccaccacaaa	cgaccccatc	Cadacccccd	gatttcccgt	tggggccacc	1440
10	accegecagt	gcctcagtct	cgtctgcgtc	ctcctcccggg	gaageggegg	tgacggccgc	1500
	gtacaaagcg	gaggggggg	cgccccagca	agaccatta	ccgaccctgg	agtgcatcct	1560
	accadacaca	agcggctgcc	tgctcccgcg	gggcccgttc	gegeegeege	cctgcaaggc	1620
	caccaccacc	adadcadccc	ccacactcta	coctagaeta	cccccacct	ccgcctctgc	1680
	gctcggctac	Caddecdeed	ccgcgctcta	gggggtggg	ggcctcaacg	ggctcccgca	1740
15	caactaccto	aggccggatt	tgctcaagga	gggccrgccg	caggictacc	cgccctatct	1800
	acctcagaag	atttotttaa	cagaagccag	tanagageeea	caatacagct	tcgagtcatt	1860
	tacctataga	agctgtaagg	tcttctttaa	cyaaycatca	ggctgtcatt	atggtgtcct	1920
	tacctgtggg	agaaatgact	acateattaa	tassatass	gaagggcagc	acaactactt	1980
	tcgccttaga	agatactata	gcatcgttga	caaaacccgc	agaaaaaact	gcccagcatg	2040
20	caataaagtc	agagttgtg	aggctggcat	ggcccctgga	ggtcgaaaat	ttaaaaagtt	2100
	Éccaaatgaa	agageegega	gagcactgga	cyclyctyct	ctcccacage	cattgggcgt	2160
	tccaaatgaa gttgattcca	ccactgatca	acctattaat	accoacttt	tcaccaggtc	aagacataca	2220
	acatgacaac	acaaaacctg	acctgttaat	gagcattgaa	ccagatgtga	tctatgcagg	2280
	Cdadaddcaa	cttctttcag	acacctccag	atataaata	acaagtctta	atcaactagg	2340
25	acatattoat	gaccagataa	tagtcaagtg	gictaaatca	ttgccaggtt	ttcgaaactt	2400
	tctaggatgg	agatectaea	ctctcattca	grattettgg	atgagettaa	tggtgtttgg	2460
		agaccccaca	aacacacacaa	Luuucaaara	CTGTatttta	0000+00+	2522
		gaacaqcaaa	Luaaauaatt	al Carterat	Trattatas.		0500
	.,,,,,,,,,,,	caggagateg	Luadyctica	auttauccaa	gaagagttcc	+ 0+ 0+ 0+ 00 0	2640
30		CCCCCCCCCC	Caattttt	QQAAQQQCTA	CCAACECAAA		2700
	ggagacgagg	Luadyulada	LLauauaucr	Catcaaddca	21100111		07.0
	-99-99	ageteacage	utttttatca	acttacaaaa	CttCttdata	20++20-4	2820
		Caacttcatt	LULACLUCLL	gaaracarrr	atcoagtoo	~~~~~+	2000
	-gergaacee	ccayaaatya	Lycciquage	tattgctgca	caattaccca	agatattggc	2940
25	agggatggtg	aaaccccttc	tettteataa				2970
35		•					

INTERNATIONAL SEARCH REPORT

ational Application No PCT/EP 00/01368

a. classification of subject matter IPC 7 C12N15/12 C12N15/57 C12N15/67 C12N15/85 C12N9/64 C07K14/72 C12Q1/68 A61K48/00 According to International Patent Classification (IPC) or to both national classification and IPC Minimum documentation searched (classification system followed by classification symbols) C12N C07K C12Q A61K Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Electronic data base consulted during the international search (name of data base and, where practical, search terms used) C. DOCUMENTS CONSIDERED TO BE RELEVANT Category : Citation of document, with indication, where appropriate, of the relevant passages Relevant to claim No. WO 94 28150 A (UNIV MCGILL) X 1,2,6,7, 8 December 1994 (1994-12-08) 11,29,30 page 5, line 1 - line 11 3-5.8.9page 6, line 34 -page 7, line 10 page 6, line 24 - line 28 page 10, line 20 - line 25 page 14, line 14 - line 19 claims 1-11 χ Further documents are listed in the continuation of box C. X Patent family members are listed in annex. Special categories of cited documents : T" later document published after the international filing date or priority date and not in conflict with the application but bited to understand the principle or theory underlying the "A" document defining the general state of the art which is not considered to be of particular relevance invention "E" earlier document but published on or after the international "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another document of particular relevance; the claimed invention citation or other special reason (as specified) cannot be considered to involve an inventive step when the document is combined with one or more other, such docu-'O" document referring to an oral disclosure, use, exhibition or ments, such combination being obvious to a person skilled document published prior to the international filing date but later than the priority date claimed "\$" document member of the same patent family Date of the actual completion of the international search Date of mailing of the international search report 6 June 2000 26/06/2000 Name and mailing address of the ISA Authorized officer European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040. Tx. 31 651 epo nl. Fax: (+31-70) 340-3016

Form PCT ISA-210 (second sheet) (July 1992)

Hornig, H

INTERNATIONAL SEARCH REPORT

Int. stional Application No PCT/EP 00/01368

2.40		PCT/EP 00	/01368
	ation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages		Relevant to claim No.
X	V. BOONYARATANAKORNKIT ET AL.: "High-mobility group chromatin proteins 1 and 2 functionally interact with steroid hormone receptors to enhance their DNA binding in vitro and transcriptional activity in mamalian cells" MOL. CELL. BIOL., vol. 18, no. 8, August 1998 (1998-08), pages 4471-4487, XPO02139580 ASM WASHINGTON, DC,US cited in the application the whole document		1,2,7
X	WO 94 17182 A (RES INST OF THE PALO ALTO MEDI ;LEAVITT JOHN C (US)) 4 August 1994 (1994-08-04) page 16, line 30 - line 36 page 17, line 1 - line 3; claims 1-16		1,2,6,7, 11,29,30
X	WO 93 20218 A (CONNAUGHT LAB ;FILMUS JORGE (CA); KLEIN MICHEL (CA)) 14 October 1993 (1993-10-14) the whole document	· · · · .	1,2,6,11
Υ .	WO 94 29471 A (GENETIC THERAPY INC) 22 December 1994 (1994-12-22) the whole document		3-5,8,9
A	WO 93 23431 A (BAYLOR COLLEGE MEDICINE) 25 November 1993 (1993-11-25) cited in the application the whole document		·
A	BEATO M ET AL: "Transcriptional regulation by steroid hormones" STEROIDS: STRUCTURE, FUNCTION, AND REGULATION, US, ELSEVIER SCIENCE PUBLISHERS, NEW YORK, NY, vol. 61, no. 4, 1 April 1996 (1996-04-01),		
	pages 240-251, XP004026583 ISSN: 0039-128X the whole document		
A	BEATO M: "GENE REGULATION BY STEROID HORMONES" CELL,US,CELL PRESS, CAMBRIDGE, NA, vol. 56, no. 3, 10 February 1989 (1989-02-10), pages 335-344, XP000051659 ISSN: 0092-8674 the whole document		
	-/- -		
	·		

.

In. ational Application No
PCT/EP 00/01368

		PCT/EP 00	0/01368
C.(Continu	etion) DOCUMENTS CONSIDERED TO BE RELEVANT Citation of document, with indication, where appropriate, of the relevant passages		
			Relevant to claim No.
Α	KURACHI S. ET AL: "Regulatory mechanism of human factor IX gene: Protein binding at the Leyden-specific region." BIOCHEMISTRY, (1994) 33/6 (1580-1591)., XP002139581 the whole document		
A	CROSSLEY M. ET AL: "Recovery from hemophilia B Leyden: An androgen-responsive element in the factor IX promoter." SCIENCE, (1992) 257/5068 (377-379)., XP002139582 the whole document		
	30 1		

INTERNATIONAL SEARCH REPORT

Information on patent family members

PCT/EP 00/01368

			·		PCT/EP	00/01368	
Patent document cited in search report			Publication date	Patent family member(s)		Publication date	
WO 94	428150 	Α	08-12-1994	US 55124 AU 67918		30-04-1996 20-12-1994	
WO 94	417182	A	04-08-1994	AU 🔄 608769	94 A	15-08-1994	
WO 93	320218	A .	14-10-1993	AU 388339 BR 930616 EP 063394 FI 94449 JP 270198 JP 750145 NO 94365 US 555902	67 A 41 A 51 A 83 B 56 T 10 A	08-11-1993 13-01-1998 18-01-1995 26-09-1994 21-01-1998 16-02-1995 30-11-1994 24-09-1996	
WO 94	129471	Α	22-12-1994	EP 071028 JP 851142 US 593593	23 T	08-05-1996 03-12-1996 10-08-1999	•
WO 93	323431	Α	25-11-1993	US 536479 AU 68505 AU 424179 AU 606519 CA 213564	54 B 93 A 98 A	15-11-1994 15-01-1998 13-12-1993 02-07-1998 25-11-1993	
				EP 074512 JP 750969 US 593593 US 587453	21 A 94 T 84 A	04-12-1996 26-10-1995 10-08-1999 23-02-1999	